

PUBLICATIONS - K.S. MURRAY (Nov. 2007)

341. Redox-Active Complexes Containing Group 8 Metal Centers Linked by C2 Bridges. M. I. Bruce, K. Costuas, B. G. Ellis, J-F. Halet, P. J. Low, B. Moubaraki, K. S. Murray, N. Ouddaie, G. J. Perkins, B. W. Skelton, A. H. White, *Organometallics*, 2007, **26**, 3735-3745.
340. Coordination chemistry of 1,4-bis(carboxymethyl)-1,4,7-triazacyclononane: Synthesis and characterization of mononuclear and binuclear μ -oxo-bridged iron(III) complexes, and a 1D-helical copper(II) chain. U. Kreher, Ute, M. T. W. Hearn, B. Moubaraki, K. S. Murray, L. Spiccia, *Polyhedron*, 2007, **26**, 3205-3216.
339. Solvothermal vs bench-top reactions: control over the formation of discrete complexes and coordination polymers. A. S. R. Chesman, D. R. Turner, D. J. Price, B. Moubaraki, K. S. Murray, G. B. Deacon, S. R. Batten, *Chem. Comm.* 2007, 3541-3543.
338. Anion-solvent dependence of bistability in a family of meridional N-donor ligand containing iron(II) spin crossover complexes. B. A. Leita, S. M. Neville, G. J. Halder, B. Moubaraki, C. J. Kepert, J.-F. Létard, K. S. Murray, *Inorg. Chem.* 2007, **46**, 8784-8795.
337. Structure, Magnetism and Photomagnetism of Mixed - Ligand Tris(pyrazolyl)methane Iron(II) Spin Crossover Compounds. B. Moubaraki, B. A. Leita, S. R. Batten, P. Jensen, K. S. Murray, J. P. Smith, J. D. Cashion, C. J. Kepert, G. J. Halder, J.-F. Létard, *Dalton Trans.* 2007, 4413-4426.
336. Magnetochemistry of molecular materials: Part 2. K. S. Murray, *Chem. Aust.* 2007, 74, Mar. pp. 4-5 (invited article)
335. Magnetochemistry of molecular materials: Part 1. K. S. Murray, *Chem. Aust.* 2007, 74, Jan/Feb. pp. 5-7. (invited article)
334. A thermal spin transition in a nanoporous iron(II) coordination framework material. S. M. Neville, B. Moubaraki, K.S. Murray, C. J. Kepert, *Angew. Chem. Int. Ed.* 2007, **46**, 2059-2062.
333. Spin-crossover studies on a series of 1D chain and dinuclear iron(II) triazine-dipyridylamine compounds. S. M. Neville, B. A. Leita, D. A. Offermann, M. B. Duriska, B. Moubaraki, K. W. Chapman, G. J. Halder, K. S. Murray, *Eur. J. Inorg. Chem.*, 2007, 1073-1085. **(invited front cover)**
332. Incorporation of a tripodal ligand with a (N,O,O)-donor set into a new family of nickel and cobalt spin clusters. K. G. Alley, R. Bircher, H. A. Güdel, B. Moubaraki, K. S. Murray, B. F. Abrahams and C. Boskovic, *Polyhedron*, 2007, **26**, 369-377.
331. Stabilisation of heterobimetallic networks by crown ethers and hydrogen-bonding in $[\text{Ni}(\text{H}_2\text{O})_6][\text{Cu}_3\text{Cl}_8(\text{H}_2\text{O})_2] \cdot (15\text{-crown-5})_2 \cdot 2\text{H}_2\text{O}$: Structure and magnetism, P. C. Junk, B. Moubaraki and K. S. Murray, *Polyhedron*, 2007, **26**, 237-243..
330. Synthesis, crystal structure and magnetism of a single molecule magnet $[\text{Mn}_{16}\text{O}_{16}(\text{OMe})_6(\text{OAc})_{16}(\text{MeOH})_3(\text{H}_2\text{O}_3)] \cdot 6\text{H}_2\text{O}$, and of a mixed bridge 1D chain $[\text{Mn}(\mu\text{-OMe})(\mu\text{-OAc})_2]_n$, D. J. Price, S. R. Batten, B. Moubaraki and K. S. Murray, *Polyhedron*, 2007, **26**, 305-317.

329. The magnetic and structural elucidation of 3,5-bis(2-pyridyl)-1,2,4-triazolate – bridged dinuclear iron(II) spin crossover compounds. C.J. Schneider, J.D. Cashion, B. Moubaraki, S.M. Neville, S. R. Batten, D. R. Turner and K.S. Murray, *Polyhedron*, 2007, **26**, 1764-1772.
328. Dicopper(II) complexes of a new pyrazolate-containing Schiff-base macrocycle and related acyclic ligand. D. de Geest, A. Noble, B. Moubaraki, K.S. Murray and S. Brooker, *Dalton Trans.*, 2007, 467-475
327. [2 x 2] nickel(II) grid and a copper(II) square result from differing binding modes of a pyrazine-based diamide ligand. J. Klingele, K.A. Hunter, B. Moubaraki, K.S. Murray, K.J. Berry, J.F. Boas, J.R. Pilbrow, G.B. Jameson, P.D.W. Boyd and S. Brooker, *Dalton Trans.*, 2007, 633-645. **Hot article.**
326. Probing the dinucleating behavior of a bis-bidentate ligand: synthesis and characterization of some di- and mononuclear cobalt(II), nickel(II), copper(II) and zinc(II) complexes of 3,5-di(2-pyridyl)-4-(1H-pyrrol-1-yl)-4H-1,2,4-triazole. Klingele, M. H., Boyd, P. D. W., Moubaraki, B., Murray, K. S., Brooker, S. *Eur. J. Inorg. Chem.* 2006, 573-589.
325. Oxalato-bridged dinuclear copper(II) complexes of N-alkylated derivatives of 1,4,7-triazacyclononane: synthesis, X-ray crystal structures and magnetic properties. M. J. Belouseff, B. Graham, B. Moubaraki, K. S. Murray and L. Spiccia, *Eur. J. Inorg. Chem.*, 2006, 4872-4878.
324. Synthesis, X-ray crystal structures, magnetism, and phosphate ester cleavage properties of Copper(II) complexes of N-substituted derivatives of 1,4,7-triazacyclononane. M. J. Belouseff, M. B. Duriska, B. Graham, S. R. Batten, B. Moubaraki, K. S. Murray and L. Spiccia, *Inorg. Chem.*, 2006, **45**, 3746-3755.
323. New discrete and polymeric supramolecular architectures derived from dinuclear (bis- β -diketonato)copper(II) metallocycles, J. K. Clegg, K. Gloe, M. J. Hayter, O. Kataeva, L. F. Lindoy, B. Moubaraki, J. C. McMurtrie, K. S. Murray and D. Schilter, *Dalton Trans*, 2006, 3977-3984.
322. 1D chain structures of $M(dca)_2(phen)(H_2O).MeOH$, $M=Mn, Fe, Co, Ni, Zn, Cd$, $dca = dicyanamide, N(CN)_2^-$, $phen = 4,7$ -phenanthroline. D. S. Tonzing, S. R. Batten and K. S. Murray, *J. Mol. Struct.*, 2006, **796**, 63-68.
321. Mixed-valent cobalt spin clusters: a hexanuclear complex and a one-dimensional coordination polymer comprised of alternating hepta- and mononuclear fragments. K. G. Alley, R. Bircher, O. Waldmann, S. T. Oschenbein, H. A. Güdel, B. Moubaraki, K. S. Murray, F. Fernandez-Alonso, B. F. Abrahams and C. Boskovic, *Inorg. Chem.* 2006, **45**, 8950-8957.
320. Identification of the dinuclear and tetranuclear air-oxidised products derived from labile phenolate-bridged dimanganese(II) pyridyl-chelate compounds, F. B. Larsen, A. Boisen, K. J. Berry, B. Moubaraki, K. S. Murray, V. McKee, R. C. Scarrow and C. J. McKenzie, *Eur. J. Inorg. Chem.*, 2006, 3841-3852.
319. Citrate, in collaboration with a guanidinium ion, as a generator of cubane-like complexes with a range of metal ions: synthesis, structures, and magnetic properties of $[C(NH_2)_3]_8[(M^{II})_4(cit)_4].8H_2O$ ($M = Mg, Mn, Fe, Co, Ni, \text{ and } Zn$; $cit = citrate$), T. A. Hudson, K. J. Berry, B. Moubaraki, K.S. Murray and R. Robson, *Inorg. Chem.*, 2006, **45**, 3549-3556.

318. Structural and magnetic resolution of a two-step full spin-crossover transition in a dinuclear iron(II) pyridyl-bridged compound. J.J.M. Amoore, C.J. Kepert, J.D. Cashion, B. Moubaraki, S.M. Neville and K.S. Murray, *Chem. Eur. J.* 2006, **12**, 8220-8227. **Invited front cover of the issue.**
317. Heptanuclear iron(III) triethanolamine clusters exhibiting 'millennium dome'-like topologies and an octanuclear analogue with ground spin states of $S = 5/2$ and 0, respectively. L.F. Jones, P. Jensen, B. Moubaraki, K.J. Berry, J.F. Boas, J.R. Pilbrow, and K.S. Murray, *J. Mater. Chem.* 2006, **16**, 2690-2697. **Highlighted as a "Hot" paper in J. Mater. Chem. in the Special Issue on 'Molecular Magnetic Materials'; Edited by E. Coronado and D. Gatteschi.**
316. New mixed-valence $Mn^{II}Mn^{III}_2$ clusters exhibiting an unprecedented $Mn^{II/III}$ oxidation state distribution in their cores. L.M. Wittick, L.F. Jones, P. Jensen, B. Moubaraki, L. Spiccia, K.J. Berry and K.S. Murray, *Dalton Trans.*, 2006, 1534-1543. **Highlighted in Chemical Science; RSC Publishing, issue 3, 2006.**
315. Structure and magnetic properties of polynuclear chloro- and hydroxo-bridged copper(II) complexes formed by a tetramacrocyclic derivative of 1,4,7-triazacyclononane. A.R. Battle, B. Graham, L. Spiccia, B. Moubaraki, K.S. Murray, B.W. Skelton and A.H. White, *Inorg. Chim. Acta*, 2006, **359**, 289-297.
314. Benzotriazole based 1-D, 2-D and 3-D metal dicyanamide and tricyanomethanide coordination networks. L.F. Jones, L. O'Dea, D.A. Offerman, P. Jensen, B. Moubaraki, K.S. Murray, *Polyhedron*, 2006, **25**, 360-372.
313. The role of diisopropanolamine (dipaH₃) in cluster dimerisation and polymerisation: from spin frustrated $S = 5Fe^{III}_6$ clusters to the novel 1-D covalent polymer of mixed valence $[Co^{II}_3Co^{III}]$ tetramers. L.F. Jones, P. Jensen, B. Moubaraki, J.D. Cashion, K.J. Berry and K.S. Murray, *Dalton Trans.*, 2005, 3344-3352.
312. Copper(II) chloride complexes with multimodal ligands based on the cyclotriphosphazene platform. E.W. Ainscough, A.M. Brodie C.V. Depree, B. Moubaraki, K.S. Murray and C.A. Otter, *Dalton Trans.*, 2005, 3337-3343.
311. Synthesis and some first-row transition-metal complexes of the 1,2,4-triazole-based bis(terdentate) ligands TsPMAT and PMAT. M.H. Klingele, B. Moubaraki, K.S. Murray and S. Brooker, *Chem. Eur. J.*, 2005, **11**, 6962-6973.
310. Square planar versus tetrahedral NiS_4 cores in the Coordination spheres of (HMB)Ru(II) and $Cp^*Ru(III)$ and a related CuS_4 complex. Synthetic, single-crystal x-ray diffraction, and magnetic studies $\{HMB = \eta-C_6Me_6 \text{ and } Cp^* = \eta^5-C_5Me_5\}$. R.Y.C. Shin, M.E. Teo, G.K. Tan, L.L. Koh, J.J. Vittal, L.Y. Goh, K.S. Murray and M. Boubaraki, *Organometallics*, 2005, **24**, 4265-4273.
309. Synthetic, EPR spectroscopic, magnetic and X-ray crystallographic structural studies on copper(II) complexes of the tridentate N₂S donor ligand formed from 6-methyl-2-formylpyridine and S-methyldithiocarbamate. A.M. Ali, A.H. Mirza, R.J. Fereday, R.J. Butcher, J.M. Fuller, S.C. Drew, L.R. Gahan, G.R. Hanson, B. Moubaraki and K.S. Murray, *Inorg. Chim. Acta*, 2005, **358**, 3937-3948.
308. High-nuclearity mixed-chelate ferric complexes with a new family of polynuclear precursors. C. Boskovic, H.A. Güdel, A. Neels, W. Wernsdorfer, B. Moubaraki and K.S. Murray, *Inorg. Chem.*, 2005, **44**, 3181-3189.

307. Self assembly of the octanuclear cluster $[\text{Cu}_8(\text{OH})_{10}(\text{NH}_2(\text{CH}_2)_2\text{CH}_3)_{12}]^{6+}$ and the one dimensional N-propylcarbamato-linked coordination polymer $\{[\text{Cu}(\text{O}_2\text{CNH}(\text{CH}_2)_2\text{CH}_3)_3](\text{ClO}_4)\}_n$. F. Bramsen, A. D. Bond, C. J. McKenzie, B. Moubaraki, K. S. Murray, *Chem. Eur. J.* 2005, **11**, 825-831. **Front cover.**
306. First complexes of a 4-alkyl-3,5-di(2-pyridyl)-4*H*-1,2,4-triazole: Synthesis, x-ray crystal structures and magnetic properties of dinuclear cobalt (II), nickel(II) and copper(II) complexes of 4-isobutyl-3,5-di(2-pyridyl)-4*H*-1,2,4-triazole, M.H. Klingele, P.D.W. Boyd, B. Moubaraki, K.S. Murray and S. Brooker, *Eur. J. Inorg. Chem.* 2005, 910-918.
305. The role of acid in the formation of hydrogen-bonded networks featuring 4,4'-dicarboxy-2,2'-bipyridine (H_2dcbp): Synthesis, structural and magnetic characterisation of $\{[\text{Cu}(\text{H}_2\text{dcbp})\text{Cl}_2]\cdot\text{H}_2\text{O}\}_2$ and $[\text{Cu}(\text{H}_2\text{dcbp})(\text{NO}_3)_2(\text{H}_2\text{O})]$, E. Tynan, P. Jensen, A.C. Lees, B. Moubaraki, K.S. Murray and P.E. Kruger, *Cryst. Eng. Comm.*, 2005, **7**, 90-95.
304. Dinuclear copper(II) complexes of two homologous pyrazine-based bis-terdentate diamide ligands, J. Klingele (nee Hausmann), B. Moubaraki, K.S. Murray, J. F. Boas and S. Brooker, *Eur. J. Inorg. Chem.*, 2005, 1530-1541.
303. Spin-crossover in dimeric hydrogen-bonded iron(II) 2-(pyrazolyl)-pyridine and 2-(imidazolyl)-pyridine complexes, B.A. Leita, B. Moubaraki, K.S. Murray and J.P. Smith, *Polyhedron*, 2005, **24**, 2165-2172.
302. The first X-ray crystal structure determination of a dinuclear complex trapped in the [low spin-high spin] state: $[\text{Fe}^{\text{II}}_2(\text{PMAT})_2](\text{BF}_4)_4\cdot\text{DMF}$, M.H. Klingele, B. Moubaraki, J.D. Cashion, K.S. Murray and S. Brooker, *Chem. Comm.*, 2005, 987-989. **Front cover.**
301. Hybrid materials containing organometallic cations and 3-D anionic metal dicyanamide networks of type $[\text{Cp}^*_2\text{M}][\text{M}'(\text{dca})_3]$, P.M. Van der Werff, E. Martinez-Ferrero, S.R Batten, P. Jensen, C. Ruiz-Pérez, M. Almeida, J. C. Waerenborgh, J.D. Cashion, B. Moubaraki, J.M. Martínez-Agudo, E. Coronado and K.S. Murray, *Dalton Trans.*, 2005, 285-290.
300. Designing dinuclear iron(II) spin crossover complexes. Structure and magnetism of dinitrile-, dicyanoamido-, tricyanomethanido-, bipyrimidine- and tetrazine-bridged compounds, S.R. Batten, J. Bjernemose, P. Jensen, B.A. Leita, K.S. Murray, B. Moubaraki, J.P. Smith and H. Toftlund, *Dalton Trans.*, 2004, 3370-3375.
299. The ligand, the metal and the holey host: synthesis, structural and magnetic characterisation of Co(II), Ni(II) and Mn(II) organic frameworks incorporating 4,4'-dicarboxy-2,2'-bipyridine, E. Tynan, P. Jensen, N.R. Kelly, P. E. Kruger, A. C. Lees, B. Moubaraki and K.S. Murray, *Dalton Trans.*, 2004, 3440-3447.
298. Coordination polymers of dicyanamide and methylpyrazine; Synthesis, structures and magnetic properties, A.M. Kutasi, A.R. Harris, S.R. Batten, B. Moubaraki and K.S. Murray, *Crystal Growth and Design* 2004, **4**, 605-610.
297. Structure and magnetism of 3D-anionic metal dicyanamide $(\text{MePh}_3\text{P})[\text{M}(\text{dca})_3]$ ($\text{M} = \text{Fe}, \text{Co}, \text{Ni}$) and $(\text{EtPh}_3\text{P})[\text{M}(\text{dca})_3]$ ($\text{M} = \text{Mn}, \text{Co}, \text{Ni}$), P.M. Van der Werff, S.R. Batten, P. J Jensen, B. Moubaraki and K.S. Murray, *Crystal Growth and Design* 2004, **4**, 503-508.
296. Triangles and tetrahedra: metal directed self-assembly of metallo-supramolecular structures incorporating bis- β -diketonato ligands, J.K. Clegg, L.F. Lindoy, B. Moubaraki, K.S. Murray and J.C. McMurtrie, *Dalton Trans.*, 2004, 2417-2423.

295. Coordinative flexibility of 1,2-bis[triazacyclonon-1-yl]propan-2-ol in mononuclear and binuclear Ni(II) complexes, M. Warren, A.R. Battle, B. Moubaraki, K.S. Murray, L. Spiccia, B.W. Skelton and A.H. White, *Dalton Trans.*, 2004, 2309-2313.
294. Dimetallic complexes of acyclic pyridine-armed ligands derived from 3,6-diformylpyridazine, P.G. Plieger, A.J. Downard, B. Moubaraki, K.S. Murray and S. Brooker, *Dalton Trans.*, 2004, 2157-2165.
293. Hexacyanometallates as templates for discrete pentanuclear and heptanuclear bimetallic clusters, L. Spiccia, K.S. Murray and J.F. Young, *Inorganic Synthesis* (Ed. J.R. Shapley) 2004, **34**, 133-141.
292. Synthesis, structure and magnetism of new single molecule magnets composed of Mn(II)₂Mn(III)₂ alkoxo-carboxylate bridged clusters capped by triethanolamine ligands, L. Wittick, K.S. Murray, B. Moubaraki, S.R. Batten, L. Spiccia and K.J. Berry, *Dalton Trans.*, 2004, 1003-1011.
291. Cooperativity in spin-crossover systems. Memory, magnetism and microporosity, K. S. Murray and C. J. Kepert, *Topics Curr. Chem.*, (Eds. P. Gütllich, H.A. Goodwin), 2004, **233**, 195-228.
290. Structure and magnetism of a new pyrazolate bridged iron(II) spin crossover complex displaying a single HS-HS to LS-LS transition, B.A. Leita, B. Moubaraki, K.S. Murray, J.P. Smith and J.D. Cashion, *Chem. Commun.*, 2004, 156-157.
289. Syntheses, structural properties and catecholase activity of copper(II) complexes with reduced Schiff base *N*-(2-hydroxybenzyl)-amino acids, C.-T. Yang, M. Vetrichelvan, X. Yang, B. Moubaraki, K.S. Murray and J.J. Vittal, *Dalton Trans.*, 2004, 113-121.
288. Structure and magnetism of coordination polymers containing dicyanamide and tricyanomethanide, S.R. Batten and K.S. Murray, *Coordination Chemistry Reviews*, 2003, **246**, 103-130.
287. Synthesis, structure and magnetism of $\{[\text{Mn}(\mu\text{-OH})(\mu\text{-OAc})_2]\cdot\text{HOAc}\cdot\text{H}_2\text{O}\}_n$ and the facilitation of long-range magnetic order through hydrogen bonding, D.J. Price, S.R. Batten, B. Moubaraki, K.S. Murray, *Polyhedron*, 2003, **22**, 2161-2167.
286. ID Manganese(III) and iron(III) coordination polymers containing Schiff-base ligands and dicyanamide, D.J. Price, S.R. Batten, B. Moubaraki and K.S. Murray, *Indian Journal of Chemistry*, 2003, **42A**, 2256-2266.
285. Binuclear Copper(II) Complexes of Xylyl-Bridged Bis(1,4,7-triazacyclononane) Ligands, F.H. Fry, L. Spiccia, P. Jensen, B. Moubaraki, K.S. Murray and E.R.T. Tiekink, *Inorg. Chem.*, 2003, **42**, 5594-5560.
284. Crystal structure of di- μ_2 -benzoato-di-[tris(pyrazolyl)methane-aqua-manganese(II)] benzoate, C₂₄H₂₂N₆MnO₅, B. Moubaraki, K.S. Murray and E.R.T. Tiekink, *Z. Kristallogr.*, 2003, **218**, 357-358.
283. Crystal structure of [bis-tris(pyrazolyl)methanecobalt(II)][chloro-bis(η^2 -benzoato)- η^1 -benzoato-manganate(II)]hydrate, Co(C₁₀H₁₀N₆)Mn(C₇H₅O₂)₃Cl.H₂O, B. Moubaraki, K.S. Murray and E.R.T. Tiekink, *Z. Kristallogr.*, 2003, **218**, 354-356.

282. Crystal structure of [bis-tris(pyrazolyl)methanecopper(II)][chloro-bis(η^2 -benzoato)- η^1 -benzoato-manganate(II)]hydrate, $\text{Cu}(\text{C}_{10}\text{H}_{10}\text{N}_6)\text{Mn}(\text{C}_7\text{H}_5\text{O}_2)_3\text{Cl}\cdot\text{H}_2\text{O}$, B. Moubaraki, K.S. Murray and E.R.T. Tiekink, *Z. Kristallogr.*, 2003, **218**, 351-353.
281. Crystal structure of μ_2 -oxo-bis- μ_2 -formato-bis(tris(pyrazolyl)-methane)diiron(III) perchlorate, $\text{C}_{22}\text{H}_{22}\text{Cl}_2\text{Fe}_2\text{N}_{12}\text{O}_{13}$, B. Moubaraki, K.S. Murray and E.R.T. Tiekink, *Z. Kristallogr.*, 2003, **218**, 349-350.
280. Crystal structure of *N*-(2-aminoketyl)pyridyl(salicylaldiminato)-copper(II) perchlorate dimer, $[\text{Cu}(\text{C}_{14}\text{H}_{13}\text{N}_2\text{O})_2\text{ClO}_4]_2$, M. Burns, B. Moubaraki, K.S. Murray and E.R.T. Tiekink, *Z. Kristallogr.*, 2003, **218**, 347-348.
279. Crystal structure of *trans*-diaqua-di(bis(pyrazolylmethane)copper(II) dinitrate, $\text{Cu}(\text{C}_7\text{H}_{10}\text{N}_4)_2(\text{OH}_2)_2(\text{NO}_3)_2$, K.A. Van Langenberg, B. Moubaraki, K.S. Murray and E.R.T. Tiekink, *Z. Kristallogr.*, 2003, **218**, 345-346.
278. Coordination Algorithms Control Molecular Architecture: $[\text{Cu}^{\text{I}}_4(\text{L}2)_4]^{4+}$ Grid Complex Versus $[\text{M}^{\text{II}}_2(\text{L}2)_2\text{X}_4]^{y+}$ Side-By-Side Complexes (M = Mn, Co, Ni, Zn; X = Solvent or Anion) and $[\text{Fe}^{\text{II}}(\text{L}2)_3][\text{Cl}_3\text{Fe}^{\text{III}}\text{OFe}^{\text{III}}\text{Cl}_3]$, Y. Lan, D.K. Kennepohl, B. Moubaraki, K.S. Murray, J.D. Cashion, G.B. Jameson and S. Brooker, *Chem. Eur. J.*, 2003, **9**, 3772-3784.
277. Synthesis and structural characterisation of two coordination polymers (molecular ladders) incorporating $[\text{M}(\text{OAc})_2]_2$ secondary building units and 4,4'-bipyridine [M = Cu(II), Zn(II)], B. Conerney, P. Jensen, P.E. Kruger, B. Moubaraki and K.S. Murray, *CrystEngComm*, 2003, **5**, 454-458.
276. Synthesis and structural and magnetic characterisation of tetranuclear Cu(II) complexes possessing novel $[\text{Cu}_4(\mu_4\text{-PO}_4)_2(\mu_2\text{-CO}_3)]$ butterfly cores that exhibit supramolecular isomerism, R. Doyle, P.E. Kruger, B. Moubaraki, K.S. Murray and M. Nieuwenhuyzen, *Dalton Trans.*, 2003, 4230-4237.
275. Dicobalt(II) complexes of a triazolate-containing Schiff-base macrocycle: synthesis, structure and magnetism, U. Beckmann, S. Brooker, C.V. Depree, J.D. Ewing, B. Moubaraki and K.S. Murray, *J. Chem. Soc. Dalton Trans.*, 2003, 1308-1313.
274. Synthesis, characterization and properties of ternary copper(II) complexes containing reduced Schiff base *N*-(2-hydroxybenzyl)- α -amino acids and 1,10-phenanthroline, C-T. Yang, B. Moubaraki, K.S. Murray and J.J. Vittal, *J. Chem. Soc. Dalton Trans.*, 2003, 880-889.
273. Asymmetry in endogenously bridged binuclear copper(II) and zinc(II) complexes formed by 1,2-bis[1,4,7-triazacyclonon-1-yl]propan-2-ol, F.H. Fry, B. Moubaraki, K.S. Murray, L. Spiccia, M. Warren, B.W. Skelton and A.H. White, *J. Chem. Soc. Dalton Trans.*, 2003, 866-871.
272. Structure and magnetism of trinuclear and tetranuclear mixed valent manganese clusters from dicyanonitrosomethanide derived ligands, D. J. Price, S.R. Batten, K.J. Berry, B. Moubaraki and K.S. Murray, *Polyhedron*, 2003, **22**, 165-176.
271. Guest-dependent spin crossover in a nanoporous molecular framework material, G.J. Halder, C.J. Kepert, B. Moubaraki, K.S. Murray and J.D. Cashion, *Science*, 2002, **298**, 1762-1765.

270. Synthesis, crystal structures and magnetic properties of linear and bent trinuclear complexes formed by hexacyanometallates and copper(II) complexes, R.J. Parker, K.D. Lu, S.R. Batten, B. Moubaraki, K.S. Murray, L. Spiccia, J.D. Cashion, A.D. Rae, A.C. Willis, *J. Chem. Soc., Dalton Trans.*, 2002, 3723-3730.
269. Synthesis, crystal structures, and magnetic properties of first row transition metal coordination polymers containing dicyanamide and 4,4'-bipyridine, P. Jensen, S.R. Batten, B. Moubaraki and K.S. Murray, *J. Chem. Soc. Dalton Trans.*, 2002, 3712-3722.
268. Synthesis, molecular structure and magnetic properties of $\text{Fe}_6(\mu_3\text{-O})_2(\text{OPr}^i)_2(\mu\text{-OPr}^i)_6(\text{O}_2(\text{Ph})_2(\mu\text{-O}_2\text{CPh})_4$, P. Ammala, S.R. Batten, J.D. Cashion, C.M. Kepert, B. Moubaraki, K.S. Murray, L. Spiccia and B.O. West, *Inorg. Chim. Acta.*, 2002, **331**, 90-97.
267. Exchange-coupled high-spin, low-spin and spin-crossover dicobalt(II) complexes of a pyridazine-containing Schiff-base macrocycle: control of cobalt(II) spin state by choice of axial ligands, S. Brooker, D.J. de Geest, R.J. Kelly, P.G. Plieger, B. Moubaraki, K.S. Murray and G.B. Jameson, *J. Chem. Soc., Dalton Trans.*, 2002, 2080-2087.
266. Crystal structure of η^1 -benzoato-chloro-tris(1-pyrazolylmethane)-copper(II), $\text{C}_{17}\text{H}_{15}\text{ClCuN}_6\text{O}_2$, K.A. Van Langenberg, B. Moubaraki, K.S. Murray and E.R.T. Tiekink, *Z. Krystallogr.* 2002, **217**, 223-224.
265. Crystal structure of *trans*-bis-thiocyanato-di(bis(pyrazolyl)methane)-copper(II), $\text{C}_{16}\text{H}_{16}\text{CuN}_{10}\text{S}_2$, K.A. Van Langenberg, K.S. Murray and E.R.T. Tiekink, *Z. Krystallogr.* 2002, **217**, 221-222.
264. Crystal structure of bis(η^1 -benzoato)tris(1-pyrazolyl)methanecopper(II) hydrate, $\text{C}_{24}\text{H}_{22}\text{CuN}_6\text{O}_5$, B. Moubaraki, K.S. Murray and E.R.T. Tiekink, *Z. Krystallogr.* 2002, **217**, 219-220.
263. Crystal structure of *N,N'*-1,3-diamino-2- μ -propanolato-bis(salicyclidene-amino)- μ -pyrazolato-dicopper(II), $\text{C}_{20}\text{H}_{22}\text{Cu}_2\text{N}_4\text{O}_3$, S.C. Courtney, K.S. Murray and E.R.T. Tiekink, *Z. Krystallogr.* 2002, **217**, 217-218.
262. Crystal structure of *N,N'*-ethylene-bis(1,3-dimethyl-5-pyrazolethiolatoiminomethyl)-copper(II), $\text{C}_{14}\text{H}_{18}\text{CuN}_6\text{S}_2$, A. la Cour, H. Toftlund, K.S. Murray and E.R.T. Tiekink, *Z. Krystallogr.* 2002, **217**, 215-216.
261. Structure and magnetism of heptanuclear complexes formed on encapsulation of hexacyanoferrate(II) with the Mn(II) and Ni(II) complexes of 1,4-bis(2-pyridylmethyl)-1,4,7-triazacyclononane, R.J. Parker, L. Spiccia, B. Moubaraki, K.S. Murray, D.C.R. Hockless, A.D. Rae and A.C. Willis, *Inorg. Chem.*, 2002, **41**, 2489-2495.
260. Electrochemical, spectroscopic, structural and magnetic characterization of the reduced and protonated α -Dawson anions in $[\text{Fe}(\eta^5\text{-C}_5\text{Me}_5)_2]_5\text{HS}_2\text{Mo}_{18}\text{O}_{62} \cdot 3\text{HCONMe}_2 \cdot 2\text{Et}_2\text{O}$ and $[\text{NBu}_4]_5[\text{HS}_2\text{Mo}_{18}\text{O}_{62}] \cdot 2\text{H}_2\text{O}$, S. Juraja, T. Vu. P.J.S. Richardt, A.M. Bond, T.J. Cardwell, J.D. Cashion, G.D. Fallon, G. Lazarev, B. Moubaraki, K.S. Murray and A.G. Wedd, *Inorg. Chem.*, 2002, **41**, 1072-1078.
259. Structure and magnetic properties of tri- and hexa-nuclear hydroxo-bridged copper(II) complexes formed by a trimacrocyclic derivative of 1,4,7-triazacyclononane, B. Graham, L. Spiccia, G.D. Fallon, M.T.W. Hearn, F.E. Mabbs, B. Moubaraki and K.S. Murray, *J. Chem. Soc., Dalton Trans.*, 2002, 1226-1232.

258. Utilization of crown ethers to stabilize the dinuclear μ -oxo bridged iron(III) aqua ion, $[(\text{H}_2\text{O})_5\text{Fe}(\mu\text{-O})\text{Fe}(\text{OH}_2)_5]^{4+}$, P.C. Junk, B.J. McCool, B. Moubaraki, K.S. Murray, L. Spiccia, J.D. Cashion and J.W. Steed, *J. Chem. Soc., Dalton Trans.*, 2002, 1024-1029.
257. New 2D coordination polymers containing both bi- and tri-dentate dicyanamide bridges and intercalated phenazine, A.M. Kutasi, S.R. Batten, B. Moubaraki and K.S. Murray, *J. Chem. Soc., Dalton Trans.*, 2002, 819-821.
256. Synthesis, structure and magnetism of a new manganese carboxylate cluster: $[\text{Mn}_{16}\text{O}_{16}(\text{OMe})_6(\text{OAc})_{16}(\text{MeOH})_3(\text{H}_2\text{O})_3]\cdot 6\text{H}_2\text{O}$, D.J. Price, S.R. Batten, B. Moubaraki and K.S. Murray, *Chem. Comm.*, 2002, 762-763.
255. Crystal engineering with mercuric chloride, S.R. Batten, A.R. Harris, K.S. Murray and J.P. Smith, *Crystal Growth & Design*, 2002, **2**, 87-89.
254. Structural and magnetic properties of the coordination polymer $\text{Mn}(\text{dca})_2(\text{H}_2\text{O})_2\cdot 2\text{Me}_4\text{pyz}$, dca = Dicyanamide ($\text{N}(\text{CN})_2^-$), Me_4pyz = tetramethylpyrazine, A.M. Kutasi, S.R. Batten, A.R. Harris, B. Moubaraki and K.S. Murray, *Aust. J. Chem.*, 2002, **55**, 311-313.
253. Structure and magnetism of the ladder-like coordination polymer $\text{Co}_3(\text{dca})_2(\text{nic})_4(\text{H}_2\text{O})_8\cdot 2\text{H}_2\text{O}$ [dca = dicyanamide anion, $\text{N}(\text{CN})_2^-$; nic = nicotinate anion], A.M. Kutasi, S.R. Batten, A.R. Harris, B. Moubaraki and K.S. Murray, *Cryst. Eng. Comm.*, 2002, **46**, 202-204.
252. Malleable coordination networks, S.R. Batten and K.S. Murray, *Aust. J. Chem.*, 2001, **54**, 605 - 609.
251. Copper(I) tris(pyrazolyl)methane complexes and their reactivity towards dioxygen, M. Cvetkovic, S.R. Batten, B. Moubaraki, K.S. Murray, L. Spiccia, *Inorg. Chim. Acta*, 2001, **324**, 131-140.
250. A purple acid phosphatase from sweet potato contains an antiferromagnetically coupled binuclear Fe-Mn center, G. Schenk, C.L. Boutchard, L.E. Carrington, C.J. Noble, B. Moubaraki, K.S. Murray, J. De Jersey, G.R. Hanson, S. Hamilton, *J. Biol. Chem.*, 2001, **276(22)**, 19084-19088.
249. Mononitrosyl and trans-dinitrosyl complexes of phthalocyaninates of manganese and rhenium, M. Goldner, B. Geniffke, A. Franken, K.S. Murray, H. Homborg, *Z. Anorg. Allg. Chem.*, 2001, **627(5)**, 935-947.
248. Lithium intercalation in anatase-structural and magnetic considerations, V. Luca, B. Hunter, B. Moubaraki, K.S. Murray, *Chem. Mater.*, 2001, **13(3)**, 796-801.
247. Interconversion of a monomer and two coordination polymers of a copper(II)-reduced Schiff Base ligand-1,10-phenanthroline complex based on hydrogen- and coordinative-bonding, C. Tong Yang, B. Moubaraki, K.S. Murray, J.D. Ranford and J.J. Vittal, *Inorg. Chem.*, 2001, **40**, 5934-5941.
246. Structure and magnetism of anionic dicyanamidometallate extended networks of types $(\text{Ph}_4\text{As})[\text{M}^{\text{II}}(\text{dca})_3]$ and $(\text{Ph}_4\text{As})_2[\text{M}_2^{\text{II}}(\text{dca})_6(\text{H}_2\text{O})]\cdot \text{H}_2\text{O}\cdot x\text{CH}_3\text{OH}$, where dca = $\text{N}(\text{CN})_2^-$ and $\text{M}^{\text{II}} = \text{Co}$, Ni, P.M. van der Werff, S.R. Batten, P. Jensen, B. Moubaraki, K.S. Murray and E. H.-K. Tan, *Polyhedron*, 2001, **20**, 1129-1138.

245. Long-range magnetic order displayed by a bimetallic pentanuclear cluster complex $[(\text{Ni}(2,2'\text{-bipy})_2)_3(\text{Fe}(\text{CN})_6)_2]\cdot 13\text{H}_2\text{O}$ and by a Cu(II) analog, K. Van Langenberg, D.C.R. Hockless, B. Moubaraki and K.S. Murray, *Synthetic Metals*, 2001, **122**, 573-580.
244. Synthesis, structural isomerism, and magnetism of the coordination polymers $[M(\text{dca})_2\text{pyz}]$, $M=\text{Mn, Fe, Co, Ni}$ and Zn , $\text{dca} = \text{dicyanamide } (\text{N}(\text{CN})_2^-)$, and $\text{pyz} = \text{pyrazine}$, P. Jensen, S.R. Batten, B. Moubaraki and K.S. Murray, *J. Solid State Chem.*, 2001, **159**, 352-361.
243. The synthesis, structure, magnetic and Mössbauer spectral properties of an Fe-Zr carboxylate-alkoxide derivative containing an $\{\text{Fe}_2\text{Zr}_2(\mu_3\text{-O})_2\}$ core, P.S. Ammala, J.D. Cashion, C.M. Kepert, K.S. Murray, B. Moubaraki, L. Spiccia and B.O. West, *J. Chem. Soc., Dalton Trans.*, 2001, 2032-2041.
242. Doubly pyridazine-bridged macrocyclic complexes of copper in +1, +2 and mixed valent oxidation states, S. Brooker, T.C. Davidson, S.J. Hay, R.J. Kelly, D.K. Kennepohl, P.G. Plieger, B. Moubaraki, K.S. Murray, E. Bill and E. Bothe, *Coordination Chemistry Reviews*, 2001, **216-217**, 3-30.
241. Cation templation of anionic metal dicyanamide networks, P.M. van der Werff, S.R. Batten, P. Jensen, B. Moubaraki and K.S. Murray, *Inorg. Chem.*, 2001, **40**, 1718-1722.
240. Synthesis, crystal structures, magnetic properties, and EPR spectra of tetranuclear copper(II) complexes featuring pairs of "Roof-Shaped" Cu_2X_2 Dimers with hydroxide, methoxide, and azide bridges, B. Graham, M.T.W. Hearn, P.C. Junk, C.M. Kepert, F.E. Mabbs, B. Moubaraki, K.S. Murray and L. Spiccia, *Inorg. Chem.*, 2001, **40**, 1536-1543.
239. Structure and magnetic properties of a high-spin $\text{Mn}_6^{\text{II}}\text{Cr}^{\text{III}}$ cluster containing cyano bridges and Mn centres capped by pentadentate ligands, Richard J. Parker, Leone Spiccia, Kevin J. Berry, Gary D. Fallon, Boujemaa Moubaraki and Keith S. Murray, *Chem. Commun.*, 2001, 333-334.
238. Electrochemical Synthesis and Structural and Physical Characterization of One- and Two-Electron-Reduced Forms of $[\text{SMo}_{12}\text{O}_{40}]^{2-}$, T. Vu, A.M. Bond, C.R. Hockless, B. Moubaraki, K.S. Murray, G. Lazarev and A.G. Wedd, *Inorg. Chem.*, 2001, **40**, 65-72.
237. 3D networks constructed using both coordinative and hydrogen bonding: synthesis, structure and magnetic properties of $\text{M}(\text{tcm})_2(\text{H}_2\text{O})_2\cdot \text{Me}_4\text{pyz}$, $\text{M} = \text{Co, Ni}$, S.R. Batten, B.F. Hoskins, B. Moubaraki, K.S. Murray and R. Robson, *Cryst. Eng. Comm.*, 2001, **8**, 1-3.
236. Anionic metal dicyanamide networks with paramagnetic counter-cations, Stuart R Batten, Paul Jensen, Boujemaa Moubaraki and Keith S Murray, *Chem. Commun.*, 2000, 2333-2332.
235. Dicopper(II) complexes of a phenolate-hinged ditopic ligand with solvent-derived ancillary ligands bound at the potentially dimetallic exogenous site, Pia Dalgaard, Alan Hazell, Christine J McKenzie, Boujemaa Moubaraki and Keith S Murray, *Polyhedron*, 2000, **19**, 1909-1915.
234. Structures and spectra of bis-tripodal iron(II) chelates, $[\text{FeL}_2]^{2+}$, where $\text{L} = \text{tris}(\text{pyrazol-1-yl})\text{methane}$, $\text{tris}(\text{pyridin-2-yl})\text{methane}$, $\text{bis}(\text{pyrazol-1-yl})(\text{pyridin-2-yl})\text{methane}$ and $\text{tris}(\text{pyridin-2-yl})\text{-phosphine oxide}$. Magnetism and spin crossover in the $(\text{pz})_3\text{CH}$ case, Peter A Anderson, Timothy Astley, Michael A Hitchman, F. Richard Keene, Boujemaa Moubaraki, Keith S Murray, Brian W Skelton, Edward R T Tiekink, Hans Toftlund and Allan H White, *J. Chem. Soc., Dalton Trans.*, 2000, 3505-3512.

233. Tetranuclear copper(II) complexes incorporating short and long metal-metal separations: synthesis, structure and magnetism, Paul E Kruger, Boujemaa Moubaraki, Gary D Fallon and Keith S Murray, *J. Chem. Soc., Dalton Trans.*, 2000, 713-718.
232. Cyano bridged dinuclear Cu(II) complexes, Richard J Parker, Leone Spiccia, Boujemaa Moubaraki, Keith S Murray, Brian W Skelton and Allan H White, *Inorganic Chimica Acta* 2000, **300**, 922-931.
231. Tris(2,2'-bipyridyl-N,N')iron(II) diperchlorate, Stuart R. Batten, Keith S. Murray and Nathan J. Sinclair, *Acta Cryst., Sect. C*, 2000, **56**, e320-e320.
230. Phase change in tetraphenylphosphonium perchlorate, Stuart R. Batten, Alexander R. Harris and Keith S. Murray, *Acta Cryst., Sect. C*, 2000, **56**, 1394-1395.
229. Copper(I) dicyanamide coordination polymers: ladders, sheets, layers, diamond-like networks and unusual interpenetration, Stuart R. Batten, Alexander R. Harris, Paul Jensen, Keith S. Murray and Angela Ziebell, *J. Chem. Soc., Dalton Trans.*, 2000, 3829-3835.
228. Self-Penetration - a Structural Compromise between Single Networks and Interpenetration: Magnetic Properties and Crystal Structures of $[\text{Mn}(\text{dca})_2(\text{H}_2\text{O})]$ and $[\text{M}(\text{dca})(\text{tcm})]$, $\text{M} = \text{Co}, \text{Ni}, \text{Cu}$, $\text{dca} = \text{Dicyanamide}, \text{N}(\text{CN})_2^-$, $\text{tcm} = \text{Tricyanomethanide}, \text{C}(\text{CN})_3^-$, Paul Jensen, David J. Price, Stuart R. Batten, Boujemaa Moubaraki and Keith S. Murray, *Chem. Eur. J.*, 2000, **6**, 3186-3195.
227. An octanuclear Fe(III) compound featuring a new type of double butterfly iron-oxo core, P. Ammala, J.D. Cashion, C.M. Kepert, B. Moubaraki, K.S. Murray, L. Spiccia and B.O. West, *Angew. Chem. Int. Ed.* 2000, **39**, 1688-1690.
226. An alternative to interpenetration whereby nets with large windows may achieve satisfactory space filling, S.R. Batten, B.F. Hoskins, B. Moubaraki, K.S. Murray and R. Robson, *Chem. Commun.*, 2000, 1095-1096.
225. Infinite molecular tubes: structure and magnetism of $\text{M}(\text{dca})_2(\text{apym})$, $\text{M} = \text{Co}, \text{Ni}$, $\text{apym} = 2\text{-aminopyrimidine}$, $\text{dca} = \text{dicyanamide}, \text{N}(\text{CN})_2^-$, Paul Jensen, Stuart R. Batten, Boujemaa Moubaraki and Keith S. Murray, *Chem. Commun.*, 2000, 793-794.
224. X-ray Diffraction and Magnetic Studies on a Series of Isosystructural Divalent Metal Chloranilates with Zig-Zag Polymeric Chain Structures and on a Dinuclear Iron(III) Chloranilate, B.F. Abrahams, K.D. Lu, B. Moubaraki, K.S. Murray and R. Robson, *J. Chem. Soc. Dalton Trans.*, 2000, 1793-1797
223. Synthesis of $[\text{Cu}_2(\text{Me}_2[9]\text{aneN}_2\text{S})_2(\text{OH})_2](\text{PF}_6)_2$, a hydroxo-bridged copper(II) complex of N,N' -dimethyl-1-thia-4,7-diazacyclononane ($\text{Me}_2[9]\text{aneN}_2\text{S}$). X-Ray structural analysis, magnetic susceptibility, e.p.r. and visible spectroscopy. V.A. Grillo, L.R. Gahan, G.R. Hanson, T.W. Hambley, R. Stranger, B. Moubaraki, K.S. Murray, *Aust. J. Chem.*, 1999, **52**, 861-866.
222. Preparation, characterisation and structures of copper(II) complexes of an asymmetric anti-cancer drug analogue. B. Moubaraki, K.S. Murray, J.D. Ranford, J.J. Vittal, X. Wang, Y. Xu, *J. Chem. Soc. Dalton Trans.*, 1999, 3573-3578.
221. Syntheses, structures and magnetism of $\alpha\text{-Mn}(\text{dca})_2$, $[\text{Mn}(\text{dca})_2(\text{H}_2\text{O}_2)\cdot\text{H}_2\text{O}]$, $[\text{Mn}(\text{dca})_2(\text{C}_2\text{H}_5\text{OH})_2]\cdot(\text{CH}_3)_2\text{CO}$, $[\text{Fe}(\text{dca})_2(\text{CH}_3\text{OH})_2]$ and $[\text{Mn}(\text{dca})_2(\text{L})_2]$, where $\text{L} =$

- pyridine, CH₃OH or DMF and dca⁻ = dicyanamide, N(CN)₂⁻. S.R. Batten, P. Jensen, C.J. Kepert, M. Kurmoo, B. Moubaraki, K.S. Murray, D.J. Price, *J. Chem. Soc. Dalton Trans.*, 1999, 2987-2997.
- 220 Crystal structures and magnetic properties of the interpenetrating rutile-related compounds M(tcm)₂, M = octahedral, divalent metal, tcm = tricyanomethamide, C(CN)₃⁻, and the sheet structures of M(tcm)₂(EtOH)₂, M = Co, Ni. S.R. Batten, B.F. Hoskins, B. Moubaraki, K.S. Murray, R. Robson, *J. Chem. Soc. Dalton Trans.*, 1999, 2977-2986.
219. Nickel complexes with tris(2-aminoethyl)amine (tren): [Ni₃(tren)₄(H₂O)₂][Cr(ox)₃]₂.6H₂O (ox = oxalate), {[Ni₂(tren)₃][ClO₄]₄.H₂O}_n, and [Ni₂(tren)₂(aepd)][ClO₄]₄.2H₂O (aepd = N-(2-aminoethyl)pyrrolidine-3,4-diamine). Synthesis, structure and magnetism. V.M. Masters, P.V. Bernhardt, L.R. Gahan, B. Moubaraki, K.S. Murray, K.J. Berry, *J. Chem. Soc. Dalton Trans.*, 1999, 2323-2328.
218. Anti-inflammatory dinuclear copper(II) complexes with indomethacin. Synthesis, magnetism and EPR spectroscopy. Crystal structure of the N,N-dimethylformamide adduct. J.E. Weder, T.W. Hambley, B.J. Kennedy, P.A. Lay, D. MacLachlan, R. Bramley, C.D. Delfs, K.S. Murray, B. Moubaraki, B. Warwick, J.R. Biffin, H.L. Regtop, *Inorg. Chem.*, 1999, **38**, 1736-1744.
217. Supramolecular complexation of polynuclear aqua ions: a crown ether adduct of a μ-oxo-bridged iron(III) aqua dimer. P.C. Junk, B.J. McCool, B. Moubaraki, K.S. Murray, L. Spiccia, *Angew. Chem. Int. Edn. Engl.*, 1999, **38**, 2224-2226
216. Molecular magnetism in manganese dicyanamide extended network structures, K.S. Murray, S.R. Batten, B. Moubaraki, D.J. Price, R. Robson, *Mol. Cryst. Liq. Cryst.*, 1999, **335**, 313-322.
215. Synthesis, structural isomerism and magnetism of extended framework compounds of type [Cu(dca)₂(pyz)]_n, where dca = dicyanamide (N(CN)₂⁻) and pyz = pyrazine. P. Jensen, S.R. Batten, G.D. Fallon, D.C.R. Hockless, B. Moubaraki, K.S. Murray, R. Robson, *J. Solid State Chem.*, 1999, **145**, 387-393.
214. First cobalt complex to exhibit both exchange coupling and spin crossover effects: The structure and unusual magnetochemistry of (Co^{II}₂ L (NCS)₂(SCN)₂] where L = macrocyclic ligand. S. Brooker, P.S. Plieger, B. Moubaraki, K.S. Murray, *Angew. Chem. Int. Edn. Engl.*, 1999, **38**, 408-410.
213. Structural isomers of M(dca)₂ molecule-based magnets. Crystal structure of tetrahedrally coordinated sheet-like β-Zn (dca)₂ and β-Co/Zn (dca)₂, and the octahedrally coordinated rutile-like α-Co(dca)₂, where dca⁻ = dicyanamide, N(CN)₂⁻, and magnetism of β-Co (dca)₂, P. Jensen, S.R. Batten, G.D. Fallon, B. Moubaraki, K.S. Murray, D.J. Price, *Chem. Comm.*, 1999, 177-178.
212. Edge-sharing decahalodiosmate complexes in multiple oxidation states; synthesis, magnetochemistry, voltammetry and associated spectroelectrochemistry of (Ph₄P)₂[Os₂(μ-X)₂X₈] (X = Cl, Br), D.G. Humphrey, G.A. Heath and K.S. Murray, *J. Chem. Soc Dalton Trans.*, 1998, 2417-2424.
211. Iron(III) and iron(II) complexes of 1-thia-4,7-diazacyclononane([9]aneN₂S) and 1,4-dithia-7-azacyclononane([9]NS₂). X-ray structural analyses, magnetic susceptibility, Mössbauer and electronic spectroscopy, V.A. Grillo, L.R. Gahan, G.R. Hanson, R. Stranger, T.W.

- Hambley, K.S. Murray, B. Moubaraki and J.D. Cashion, *J. Chem. Soc. Dalton Trans.*, 1998, 2341-2348.
210. The ground state in tris(acetylacetonato)ruthenium(III) from low temperature single crystal magnetic properties, B.N. Figgis, P.A. Reynolds, K.S. Murray and B. Moubaraki, *Aust. J. Chem.*, 1998, **51**, 229-234.
209. Synthesis, structure and magnetism of the oxalato-bridged chromium(III) complex $[\text{NBu}^n_4][\text{Cr}_2(\text{ox})_5] \cdot 2\text{CHCl}_3$, V.M. Masters, C.A. Sharrad, P.V. Bernhardt, L.R. Gahan, B. Moubaraki and K.S. Murray, *J. Chem. Soc. Dalton Trans.*, 1998, 413-416.
208. Oxo-centred trinuclear acetate complexes containing mixed-metal clusters. Crystal structure of a chromium(III) iron(III) nickel(II) complex and magnetic properties of a dichromium(III) magnesium(II) complex, A.B. Blake, E.Sinn, A. Yavari, K.S. Murray and B. Moubaraki, *J. Chem. Soc. Dalton Trans.*, 1998, 45-49.
207. Bis(phthalocyaninato(2-))rhenium: Synthesis, properties and crystal structure, M. Göldner, H. Hückstadt, K.S. Murray, B. Moubaraki and H. Homborg, *Z. Anorg. Allgem. Chem.*, 1998, **624**, 288-294.
206. Structure and molecular magnetism of the rutile related compounds $\text{M}(\text{dca})_2$, $\text{M} = \text{Co}^{\text{II}}$, Ni^{II} , Cu^{II} , $\text{dca} = \text{dicyanamide}$, $\text{N}(\text{CN})_2^-$, S.R. Batten, P. Jensen, B. Moubaraki, K.S. Murray and R. Robson, *Chem. Comm.*, 1998, 439-440.
205. Structural and magnetic properties of an asymmetric dicopper(II) anticancer drug analogue, B. Moubaraki, K.S. Murray, J.D. Ranford, X. Wang and Y. Xu, *Chem. Comm.*, 1998, 353-354.
204. Multifield magnetisation saturation measurements on rubredoxin samples from *Clostridium Pasteurianum*, B. Moubaraki, K.S. Murray, Z. Xiao and A.G. Wedd, *J. Inorg. Biochem.*, 1997, **67**, 261.
203. Towards synthetic models for trinuclear copper active sites of ascorbate oxidase and laccase: Self-assembly, crystal structure and magnetic properties of the copper(II) complexes of tris(1,4,7-triazacyclonon-1-ylmethyl)benzene, L. Spiccia, B. Graham, M.T.W. Hearn, G. Lazarev, B. Moubaraki, K.S. Murray and E.R.T. Tiekink, *J. Chem. Soc. Dalton Trans.*, 1997, 4089-4097.
202. Structure and magnetism of a bimetallic pentanuclear cluster $[(\text{Ni}(\text{bpm})_2)_3(\text{Fe}(\text{CN})_6)_2] \cdot 7\text{H}_2\text{O}$ ($\text{bpm} = \text{bis}(1\text{-pyrazolyl})\text{methane}$). The role of the hydrogen-bonded $7\text{H}_2\text{O}$ "cluster" in long-range magnetic ordering, K. Van Langenberg, S.R. Batten, K.J. Berry, D.C.R. Hockless, B. Moubaraki and K.S. Murray, *Inorg. Chem.*, 1997, **36**, 5006-5015.
201. Mononuclear chromium(III), manganese(II) and iron(III) complexes of the pentadentate ligand 1,4-bis(2-pyridylethyl)1,4,7-triazacyclononane, G.D. Fallon, G.A. McLachlan, B. Moubaraki, K.S. Murray, L. O'Brien and L. Spiccia, *J. Chem. Soc. Dalton Trans.*, 1997, 2765-2769.
200. Synthesis, magnetism and electrochemistry of tetranuclear nickel(II) and copper(II) complexes of an unsymmetrical bis-pentadentate Schiff-base ligand, P.E. Kruger, B. Moubaraki and K.S. Murray, *Polyhedron*, 1997, **16**, 2659-2665.

199. Physical properties of polypyrrole films containing trisoxalatometallate anions and prepared from aqueous solution, N.S. Allen, K.S. Murray, R.J. Fleming, B.R. Saunders, *Synth. Met.*, 1997, **87**, 237-247.
198. A supramolecular approach to the crystallization of polynuclear aqua ions: Structure and Magnetism of an 18-Crown-6-adduct of bis(μ -hydroxo)octaaquadichromium(III) mesitylene-2-sulfonate trihydrate, A. Drljaca, D.C.R. Hockless, B. Moubaraki, K.S. Murray, L. Spiccia, *Inorg. Chem.*, 1997, **36**, 1988-1989.
197. Syntheses and characterisation of mononuclear and dinuclear iron^{III} complexes of 1,4,7-triazacyclononane-*N*-acetate (L). Crystal structures of $LFeCl_2$ and $[L_2Fe_2(\mu-O)(\mu-O_2CCH_3)](ClO_4)NaClO_4 \cdot 2H_2O$, B. Graham, L. Spiccia, B. Moubaraki, K.S. Murray, J.D. Cashion and D.C.R. Hockless, *J. Chem. Soc. Dalton Trans.*, 1997, 887-893.
196. Synthesis, structure and magnetism of iron(III) and -(II) complexes of 1-thia-4,7-diazacyclononane and *N,N'*-dimethyl-1-thia-4,7-diazacyclononane, V.A. Grillo, G.R. Hanson, T.W. Hambley, L.R. Gahan, K.S. Murray and B. Moubaraki, *J. Chem. Soc. Dalton Trans.*, 1997, 305-311.
195. μ -Carbido diporphyrinates and diphthalocyaninates of iron and ruthenium, A. Kienast, L. Galich, K.S. Murray, B. Moubaraki, G. Lazarev, J.D. Cashion and H. Homborg, *J. Porphyrins and Phthalocyanines*, 1997, **1**, 141-157.
194. Physical and spectroscopic properties of polypyrrole films containing transition-metal complexes as counteranions. B.R. Saunders, K.S. Murray, R.J. Fleming, R. Cervini, N.S. Allen, *Handbook of Organic Conductive Molecules and Polymers*, Ed. H.S. Nalwa, J. Wiley, 1997, Vol. 3, Chapter 12, pp 633-676.
193. High nuclearity ruthenium carbonyl cluster chemistry. 5. Local density functional, electronic spectroscopy, magnetic susceptibility and electron paramagnetic resonance studies on (carbido)decaruthenium carbonyl clusters, M.P. Cifuentes, M.G. Humphrey, J.E. McGrady, P.J. Smith, R. Stranger, K.S. Murray and B. Moubaraki, *J. Am. Chem. Soc.*, 1997, **119**, 2647-2655.
192. Metamagnetism in tetrachlorobis(*N*-phenylacetamidine)rhenium(IV), P.A. Reynolds, B. Moubaraki, K.S. Murray, J.W. Cable, L.M. Engelhardt and B.N. Figgis, *J. Chem. Soc. Dalton Trans.*, 1997, 263-267.
191. The crystal structure and magnetic properties of a heterodimetallic copper-vanadium complex, A. Hazell, C.J. McKenzie, B. Moubaraki and K.S. Murray, *Acta. Chem. Scand.*, 1997, **51**, 470-475.
190. Structural and solid-state examination of $Ca_4Mn_3O_{10}$ and $Sr_4Mn_3O_{10}$, H.J. Rossell, P. Goodman, S. Bulcock, R.H. March, S.J. Kennedy, T.J. White, F.J. Lincoln and K.S. Murray, *Aust. J. Chem.*, 1996, **49**, 205-217.
189. Hexacyanometalates as templates for heteropolynuclear complexes and molecular magnets: synthesis and crystal structure of $[Fe\{(CN)Cu(tpa)\}_6][ClO_4]_8 \cdot 3H_2O$, [tpa = tris(2-pyridylmethyl)amine], R.J. Parker, D.C.R. Hockless, B. Moubaraki, K.S. Murray and L. Spiccia, *Chem. Comm.*, 1996, 2879-2880.
188. First dicopper(II) complex to contain bridging macrocyclic-pyridazine units: structure, electrochemistry and magnetochemistry of $[Cu_2L(MeCN)_2(ClO_4)][ClO_4]_2$, S. Brooker, R.J. Kelly, B. Moubaraki and K.S. Murray, *Chem. Comm.*, 1996, 2579-2580.

187. Molecular magnetic materials and small clusters containing N-donor chelated metal species combined with hexacyanometallate, tris-oxalatometallate and related bridging groups, K.S. Murray, B. Moubaraki, K. van Langenberg, G.D. Fallon, D.C.R. Hockless, *ACS Symposium Series*, No. 644, Chap. 13, eds. M. Turnbull, T. Sugimoto, L.K. Thompson, 1996, pp. 201-215.
186. Binuclear chromium(III) complexes bridged by hydroxide and acetate groups, S.J. Brudenell, S.J. Crimp, J.K.E. Higgs, K.S. Murray, L. Spiccia, *Inorg. Chim. Acta*, 1996, **247**, 25-41.
185. Synthesis, x-ray structural determination and magnetic susceptibility, Mössbauer, and EPR studies of $(\text{Ph}_4\text{P})_2[\text{Fe}_2(\text{cat})_4(\text{H}_2\text{O})_2]\cdot 6\text{H}_2\text{O}$, a catecholato-bridged dimer of Iron(III), V.A. Grillo, G.R. Hanson, D. Wang, T.W. Hambley, L.R. Gahan, K.S. Murray, B. Moubaraki and C.J. Hawkins, *Inorg. Chem.*, 1996, **35**, 3568-3576.
184. Synthesis, magnetism, electrochemistry and reactivity of dinuclear vanadium(III) complexes of a macrocyclic tetraimine Schiff-base, P.E. Kruger, B. Moubaraki, K.S. Murray, *J. Chem. Soc. Dalton Trans.*, 1996, 1223-1228.
183. Spectroscopic magnetic and structural studies on the mixed-valence cuboidal clusters $[\text{Mo}_4\text{E}_4(\text{edta})_2]^{3-}$ (E=S or Se; H_4edta =ethylenediaminetetraacetic acid) and $[\text{Mo}_4\text{S}_4(\text{H}_2\text{O})_{12}]^{5+}$, W. McFarlane, M. Nasreldin, D.M. Saysell, Z-S Jia, W. Clegg, M.R.J. Elsegood, K.S. Murray, B. Moubaraki, A.G. Sykes, *J. Chem. Soc. Dalton Trans.*, 1996, 363-369.
182. Synthetic, X-ray structure, electron paramagnetic resonance and magnetic studies on the manganese(II) complexes of 1-thia-4,7-diazacyclononane ($[\text{9}]_{\text{ane}}\text{N}_2\text{S}$), L.R. Gahan, V.A. Grillo, T.W. Hambley, G.R. Hanlon, C.J. Hawkins, E.M. Proudfoot, B. Moubaraki, K.S. Murray, D. Wang, *Inorg. Chem.* 1996, **35**, 1039-1044.
181. Intramolecular ferromagnetism in a novel hexanuclear (μ -hydroxo)(μ -carbonato) copper(II) bipyridine complex. Structure of $[\text{Cu}_6(\text{bpy})(\mu\text{-CO}_3)_2(\mu\text{-OH})_2](\text{ClO}_4)_6\cdot 4\text{H}_2\text{O}$ and a dinuclear μ -carbonato complex, $[\text{Cu}_2(\text{bpy})_4(\mu\text{-CO}_3)](\text{PF}_6)_2\cdot 2\text{DMF}$, P.E. Kruger, G.D. Fallon, B. Moubaraki, K.J. Berry, K.S. Murray, *Inorg. Chem.*, 1995, **34**, 4808-4814.
180. The magnetochemistry of homo- and heterotetranuclear first-row d-block complexes, K.S. Murray, *Adv. Inorg. Chem.*, Ed. A.G. Sykes, 1995, **43**, 261-358.
179. Changes in geometries and spin states of thiolato-Schiff base NiN_2S_2 complexes containing N,N'-biphenyl backbones. H.Frydendahl, H. Toftlund, J. Becher, J.C. Dutton, K.S. Murray, L.F. Taylor, O.P. Anderson and E.R.T. Tiekink, *Inorg. Chem.*, 1995, **34**, 4467-4476.
178. Synthesis and magnetic exchange properties of linear trinuclear oxo bridged M(III)ORu(IV)OM(III) complexes (M = Fe, Cr, Mn) formed by two electron redox reactions. K.J. Berry, B. Moubaraki, K.S. Murray, P.J. Nichols, L.D. Schulz, B.O. West, *Inorg. Chem.*, 1995, **34**, 4123-4133.
177. Recent advances in the physical and spectroscopic properties of polypyrrole films, particularly those containing transition-metal complexes as counteranions, B.R. Saunders, R.J. Fleming and K.S. Murray, *Chem. Mater.*, 1995, **7**, 1082-1094.
176. Magnetic properties of binuclear manganese(II), cobalt(II), nickel(II) and copper(II) complexes of a macrocyclic ligand derived from pyridine-N-oxide, and crystal structure of

- the nickel complex. A.B. Blake, E. Sinn, A. Yavari, B. Moubaraki, K.S. Murray, *Inorg. Chim. Acta*, 1995, **229**, 281-290.
175. Bis[4-(2,6-dimethylphenyliminomethyl)-5-pyrazolethiolato]nickel(II) complexes with an unusual *trans*-planar structure. H. Toftlund, A.L. Nivorozhkin, A. la Cour, B. Adhikary, K.S. Murray, G.D. Fallon and L.E. Nivorozhkin, *Inorg. Chim. Acta*, 1995, **228**, 273-241.
174. A physical and spectroscopic study of polypyrrole films containing tetrasulfonated metallophthalocyanine and transition metal EDTA chelates. B.R. Saunders, K.S. Murray, R.J. Fleming and D.G. McCulloch, *Synth. Metals*, 1995, **69**, 363-364.
173. Crystal structure, electrospray ionization mass spectrometry, electron paramagnetic resonance, and magnetic susceptibility study of $[\text{Cu}_2(\text{ascidH}_2)(1,2\text{-}\mu\text{-CO}_3)(\text{H}_2\text{O})_2]\cdot 2\text{H}_2\text{O}$, the bis(copper(II)) complex of ascidiacyclamide (ascidH₄), a cyclic peptide isolated from the ascidian *Lissoclinum patella*, A.L. van den Brenk, K.A. Byriel, D.P. Fairlie, L.R. Gahan, G.R. Hanson, C.J. Hawkins, A. Jones, C.H.L. Kennard, B. Moubaraki and K.S. Murray, *Inorg. Chem.*, 1994, **33**, 3549-3557.
172. Crystal structure of μ -oxalato-bis[di-(2-aminoethyl)amine-copper(II)] diperchlorate monohydrate, $(\text{C}_4\text{H}_{13}\text{N}_3\text{Cu})_2(\text{C}_2\text{O}_4)(\text{ClO}_4)_2\cdot\text{H}_2\text{O}$, P.E. Kruger, K.S. Murray and E.R.T. Tiekink, *Zeitschrift für Kristallographie*, 1994, **209**, 624-625.
171. Binuclear nickel complexes with single azide bridges. Structure and properties of $[\text{Ni}_2(\text{N,N}$ -bis(2-aminoethyl)-*N'*-(2-pyridylmethyl)ethane-1,2-diamine)₂(μ -N₃)](ClO₄)₃ and $[\text{Ni}_2(1,4$ -bis(2-pyridylmethyl)-1,4,7-triazacyclononane)₂(μ -N₃)](ClO₄)₃, G.A. McLachlan, G.D. Fallon, R.L. Martin, B. Moubaraki, K.S. Murray and L. Spiccia, *Inorg. Chem.*, 1994, **33**, 4663-4668.
170. Synthesis, magnetism and electrochemistry of binuclear nickel(II) and copper(II) complexes of an unsymmetrical pentadentate Schiff-base ligand. X-ray structure of the $[\text{Ni}(\text{II})_2\text{L}(\text{pz})]$ complex, P.E. Krüger, B. Moubaraki, K.S. Murray and E.R.T. Tiekink, *J. Chem. Soc. Dalton Trans.*, 1994, 2129-2134.
169. Crystal structure of a metal-free tetranucleating macrocyclic ligand and crystal structure and magnetism of a di-manganese(III) derivative, A.J. Edwards, B.F. Hoskins, R. Robson, J.C. Wilson, K.S. Murray and B. Moubaraki, *J. Chem. Soc. Dalton Trans.*, 1994, 1837-1842.
168. Dimeric low-spin iron(III) phthalocyanines: Synthesis and properties of ferromagnetically coupled μ -oxo(acido-phthalocyaninatoiron(III)). S. Sievertsen, K.S. Murray, B. Moubaraki, K.J. Berry, Y. Korbatiéh, J.D. Cashion, J. Brown and H. Homborg, *Z. Anorg. Allgem. Chem.*, 1994, **620**, 1203-1212.
167. Preparation and characterization of high temperature superconductor/high density polyethylene composite materials, $[\text{YBa}_2\text{Cu}_3\text{O}_{7-x}/\text{HDPE}]$, A. Markiewicz, R.E. Mayer, B. Moubaraki, K.S. Murray, *Materials Res. Bull.*, 1994, **29**, 393.
166. Single-crystal low temperature magnetization studies of $[\text{Cs}_2\text{K}(\text{Mn}(\text{CN})_6)]$, A.B. Blake, B.N. Figgis, P.A. Reynolds, L.M. Engelhardt, B. Moubaraki, K.S. Murray, *J. Chem. Soc. Dalton Trans.*, 1994, 1121.
165. The synthesis, reactivity and magnetic susceptibilities of Cr(IV)porphyrin imido complexes, CrN(R), and attempts to form heterobinuclear μ -imido compounds with CrN(R)V and

- CrN(R)Fe bridges, B. Moubaraki, K.S. Murray, P.J. Nichols, S. Thomson and B.O. West, *Polyhedron*, 1994, **13**, 485-495.
164. Physical and spectroscopic studies of polypyrrole films containing transition metal EDTA chelates, B.R. Saunders, K.S. Murray, R.J. Fleming, D.G. McCulloch, L.J. Brown and J.D. Cashion, *Chem. of Materials*, 1994, **6**, 697-706.
163. High-spin manganese(II) phthalocyanines: Synthesis and spectroscopic properties of acidophthalocyaninatomanganese(II), S. Sievertsen, B. Moubaraki, K.S. Murray and H. Homborg, *Z. Anorg. Allgem. Chem.*, 1994, **620**, 682-690.
162. Physical properties of polypyrrole films containing Dicyanoaurate(I) anions, PPy-Au(CN)₂, R. Cervini, R.J. Fleming, B.J. Kennedy and K.S. Murray, *J. Mater. Chem.*, 1994, 89-97.
161. Tris-pyrazolylmethane ((pz)₃CH) Complexes of copper and iron, G.D. Fallon, S. Knight, B. Moubaraki, K.S. Murray, B.W. Skelton and A.H. White, *J. Inorg. Biochem.*, 1993, **51**, 504.
160. Design and structure of polymetallic biomodels involving short and long metal-metal separations, P.E. Kruger, G.D. Fallon, B. Moubaraki and K.S. Murray, *J. Inorg. Biochem.*, 1993, **51**, 473.
159. Magnetic properties of hexaammineruthenium(III) trithiocyanate single crystals, A.B. Blake, C.D. Delfs, L.M. Engelhardt, B.N. Figgis, P.A. Reynolds, A.H. White, B. Moubaraki and K.S. Murray, *J. Chem. Soc. Dalton Trans.*, 1993, 1417-1420.
158. The redox synthesis, structure and magnetic properties of the heterobimetallic μ -oxo compound [(py)(TPP)CrOFe(tmtaa)], P. J. Nichols, G.D. Fallon, B. Moubaraki, K.S. Murray and B.O. West, *Polyhedron*, 1993, **12**, 2205-2213.
157. The molecular structures and magnetic properties of μ -3 tetranuclear β -diketonate Cu complexes [Cu(bzac)(μ_3 -OC₂H₄OCH₃)]₄ and [Cu(μ -dbm)(μ_3 -OC₂H₄OCH₃)]₄, G. D. Fallon, B. Moubaraki, K. S. Murray, A. M. Van Den Bergen and B. O. West, *Polyhedron*, 1993, **12**, 1989-2000.
156. Physical and Spectroscopic Studies of Polypyrrole Films Containing Tetrasulfonated Metallophthalocyanine Counterions Prepared from Nonaqueous Solution, B R Sanders, K S Murray, R J Fleming and Y Korbatiéh, *Chem. of Materials*, 1993, **5**, 809-819.
155. The compressed tetragonal CuF₆⁴⁻ complex in KAlCuF₆: an angular overlap treatment of the electronic structure and magnetic exchange coupling, M Atanasov, M A Hitchman, R Hoppe, K S Murray, B Moubaraki, D Reinen and H. Stratemeier, *Inorg Chem*, 1993, **32**, 3397-3401.
154. Synthesis and crystal and molecular structure of a tetranuclear 'pair-of-dimers' nickel(II) Schiff-Base complex. Magnetism of the Cu₄ analogue. P E Kruger, G D Fallon, B Moubaraki and K S Murray, *J. Chem. Soc. Chem. Commun.*, 1992, 1726-1729.
153. Physical properties of polypyrrole films containing tetracyanonickelate(II) anions, PPy-Ni(CN)₄. R Cervini, R J Fleming and K S Murray, *J. Mater. Chem.*, 1992, 1157-1217.
152. Synthesis, X-ray crystal structures and magnetic properties of Ni₄ complexes of a macrocyclic tetranucleating ligand. A J Edwards, B F Hoskins, E H Kachab, A Markiewicz, K S Murray and R Robson, *Inorg. Chem.*, 1992, **31**, 3585-3591.

151. Single-crystal magnetic properties of $(\text{Co}(\text{NH}_3)_5(\text{OH}_2))[\text{Cr}(\text{CN})_6]$ between 2 and 300 K. P A Reynolds, C D Delfs, B N Figgis, B Moubaraki and K S Murray, *Austral J. Chem.*, 1992, **45**, 1899-1902.
150. The ground state in $\text{Cs}_2\text{KFe}(\text{CN})_6$ from single-crystal magnetic properties. P A Reynolds, C D Delfs, B N Figgis, B Moubaraki and K S Murray, *Austral. J. Chem.*, 1992, **45**, 1301-1305.
149. Magnetic exchange and zero-field splitting in the d^3 $[\text{Os}^{\text{V}}\text{Cl}_6]^-$ ion. P A Reynolds, C D Delfs, B N Figgis, M J Henderson, B Moubaraki and K S Murray, *J Chem. Soc. Dalton Trans.*, 1992, 2309-2312.
148. Structure and magnetism of cubic low-spin $4d^5$ $[\text{Ru}^{\text{III}}(\text{NH}_3)_6]\text{Br}[\text{SO}_4]$. P A Reynolds, C D Delfs, B N Figgis, L M Engelhardt, B Moubaraki and K S Murray, *J Chem. Soc. Dalton Trans.*, 1992, 2029-2032.
147. Synthesis, structure, spectral and magnetic properties of $(\mu\text{-carbonato})(\mu\text{-hydroxo})\text{-bis}[\text{N},\text{N}'\text{-bis}(2\text{-aminoethyl})\text{ethane-1,2-diamine}]$ chromium(III)] perchlorate. L Spiccia, G D Fallon, A Markiewicz, K S Murray and H Riesen, *Inorg. Chem.*, 1992, **31**, 1066-1072.
146. The reaction of the pyrophosphate ion with the antitumour complex $[\{\text{CuL}(\text{CH}_3\text{COO})_2\}]$ (HL = 2-formylpyridine thio-semicarbazone) and the single-crystal X-ray structure of $[(\text{CuL})_4\text{P}_2\text{O}_7]_n \cdot n \text{H}_2\text{O}$. E W Ainscough, A M Brodie, J D Ranford, J M Waters and K S Murray, *Inorg. Chim. Acta*, 1992, **147**, 107-115.
145. The structure and magnetism of a trimetallic $\mu\text{-oxo}$ ruthenium(IV) porphyrin-iron(III) salicylideneimine complex containing the Fe-O-Ru-O-Fe system. L D Schulz, G D Fallon, B Moubaraki, K S Murray and B O West, *J. Chem. Soc. Chem. Commun.*, 1992, 971-973.
144. Heterobinuclear oxo-bridged complexes. III. Cr(III)-O-Mo(V) compounds, synthesis and magnetic properties. R L Elliott, P E Kruger, K S Murray and B O West, *Austral. J. Chem.*, 1992, **45**, 889-896.
143. Physical properties of polypyrrole films containing sulfonated metal-phthalocyanine anions. B.R. Saunders, R.J. Fleming and K.S. Murray, *Synth. Metals*, 1992, **47**, 167-178.
142. A binuclear iron(III) complex with inequivalent metal environments. Relevance to iron oxo-proteins. G.D. Fallon, K.S. Murray, T. Quach, *J. Inorg. Biochem*, 1991, **43**, 549
141. New binucleating amide ligands and their copper(II) and manganese(III) complexes. Relevance to polymetallic bio-sites. K. Bertocello, G.D. Fallon, K.S. Murray, *J. Inorg. Biochem.* 1991, **43**, 204.
140. Manganese(III) complexes of a binucleating Schiff-base ligand based on the 1,3-diaminopropan-2-ol backbone. K. Bertocello, G. D. Fallon, K. S. Murray, E. R. T. Tiekink, *Inorg. Chem.* 1991, **30**, 3562-3568.
- 139 Conducting polypyrrole incorporating macrocyclic anions. R. Cervini, R. J. Fleming, K. S. Murray, *Polymer Materials Reprints* p. 607 (IUPAC Int. Symposium 1991, Monash Univ. Australia)
- 138 Synthesis, spectroscopic and magnetic properties of a unique binuclear $\mu\text{-oxo}$ -vanadium(IV) complex. H. Toftlund, S. Larsen, K.S.Murray, *Inorg. Chem.* 1991, **30**, 3964-3967..

137. A doubly-bridged binuclear iron(III) complex containing inequivalent metal environments. Synthesis, structure and magnetism. G.D. Fallon, A. Markiewicz, K.S. Murray and T. Quach, *J. Chem. Soc. Chem. Commun.*, 1991, 198.
136. The structure of a planar NiN₂S₂ complex of the tetraanionic form of the ligand N,N'-ethylene bis(o-mercaptobenzamide). J.C. Dutton, G.D. Fallon and K.S. Murray, *Chem. Letters* 1990, 983.
135. New binucleating amido ligands, synthesis and structures of copper(II) and manganese(III) complexes. K. Bertoncello, G.D. Fallon and K.S. Murray, *Polyhedron*, 1990, **23**, 2867.
134. Synthesis and Structure of Tetradentate Amido-phenolate Manganese(III) Complexes; Precursors for High-Valent Manganese Chelate Complexes. K. Bertoncello, G.D. Fallon, K.S. Murray, *Inorg. Chim. Acta* 1990, **174**, 57.
133. Synthesis, X-ray Crystal Structure and Magnetic Studies of a BisCopper (II) Complex of a C-spiro-binucleating linear octadentate ligand. T.M. Donlevy, L.R. Gahan, T.W. Hambley, G.R. Hanson, A. Markiewicz, K.S. Murray, I. L. Swan and S.R. Pickering, *Aust. J. Chem.* 1990, **43**, 1407.
132. Synthesis and Magnetic Properties of a μ -oxo-di(μ -acetato)-manganese (III) complex of a strapped tripodal-pyridylamine ligand N,N,N',N'-tetrakis (2-pyridylmethyl)-1,3-propanediamine. A model for the Mn₂ site of Mn-catalase enzymes. H. Toftlund, A. Markiewicz and K.S. Murray, *Acta. Chem. Scand.* 1990, **44**, 443.
131. Reaction of Dioxomolybdenum(VI) Hydroxamate and Thiohydroxamate Complexes with Organohydrazines. M.D. Fitzroy, G.D. Fallon, K.S. Murray, J.M. Frederiksen, E.R.J. Tiekink, *Inorg. Chim. Acta* 1990, **169**, 78.
130. Mössbauer Spectra, Magnetic Susceptibilities and Solution Equilibria of Heteronuclear Complexes of Cr^{III}OFe^{III} and Cr^{III}OMo^V E.N. Bakshi, R.L. Elliott, K.S. Murray, P.J. Nichols and B.O. West, *Austral. J. Chem.*, 1990, **43**, 707.
129. Synthesis, Crystal Structure and Magnetism of a Macrocyclic Binuclear Copper(II) Amino Alcohol Complex from a Metal-Directed Reaction Involving Formaldehyde and Nitroethane. P. Comba, G.A. Lawrance, T.M. Manning, A. Markiewicz, K.S. Murray, M.A. O'Leary, B.W. Skelton, A.H. White, *Aust. J. Chem.* 1990, **43**, 69.
128. Magnetism and electronic structure of a series of encapsulated first row transition metals. L.L. Martin, R.L. Martin, K.S. Murray and A.M. Sargeson, *Inorg. Chem.*, 1990, **29**, 1387.
127. A Macrocyclic Binuclear Vanadium (III) Complex with di- μ -alkoxo bridging and pentagonal-dipyramidal metal coordination. X-ray Crystal Structure of [V₂L(H₂O)₄](ClO₄)₄.2H₂O (H₂L = 1, 7, 14, 20- tetramethyl -2, 6, 15, 19 - tetra-aza-[7,7](2,6) pyridinophane-4, 7 diol) J.C. Dutton, G.D. Fallon and K.S. Murray, *J. Chem. Soc. Chem Commun.*, 1990, 64.
126. Preparation and Characterization of Copper(II) Complexes with 1,4,8,11- tetrakis (2-pyridylmethyl)-1,4,8,11-tetraazacyclotetradecane. E. Asato, H. Toftlund, S. Kida, M. Mikuriya and K.S. Murray. *Inorg. Chim. Acta.*, 1989, **165**, 207.
125. The Iron-Activated Alcohol Dehydrogenase from *Zymomonas Mobilis* : Spectroscopic and Magnetic Properties. E.N. Bakshi, P. Tse, K.S. Murray, G.R. Hanson, R.K. Scopes and A.G. Wedd, *J. Am. Chem. Soc.*, 1989, **111**, 8707.

124. Magnetism of Oxovanadium(IV) Complexes of Binucleating ligands. Oxidation to and Structure of a Mononuclear Oxovanadium(V) Complex of N,N'-(pentan-3-ol) bis(Salicylaldimine) J. C. Dutton, K.S. Murray, E.R.T. Tiekink, *Inorg. Chim. Acta*, 1989, **166**, 5.
123. Reactions of bis(acetylacetonato)dioxomolybdenum(VI) with Aroylhydrazines. Structure and ¹H NMR spectrum of 3,5-Dimethylpyrazolium β-Octamolybdate Tetrahydrate. M.D. Fitzroy, G.D. Fallon and K.S. Murray, *Inorg. Chim. Acta*, 1989, **157**, 187.
122. Electrically conducting TCNQ derivatives of copper sulphur/nitrogen chelates; Structure of a novel semiconducting complex [Cu(pdto)(TCNQ)]₂ which contains N-bonded TCNQ (pdto=1,8-di-2-pyridyl-3,6-dithiaoctane; TCNQ=7,7,8,8-tetracyanoquinodimethane). D. G. Humphrey, G. D. Fallon and K. S. Murray, *J. Chem. Soc., Chem. Comm.*, 1988, 1356-1358.
121. Exogenous bridging and non-bridging in copper(II) complexes of a binucleating 2,6-bis(N-methylpiperazino)methyl-4-chlorophenolate ligand. Crystal structures and magnetic properties of bis(μ-acetato), Dinitrito and bis (azido) Complexes. Possible Relevance to the Type 2 Depleted Laccase Active Site. K. Bertoncello, G.D. Fallon, J.H. Hodgkin, K.S. Murray, *Inorg. Chem.*, 1988, **27**, 4750-4758.
120. Iron(IV) Phthalocyanines. Mössbauer Spectral Studies of μ-Carbido-Phthalocyaninato Iron(IV) and its Axially Ligated and Oxidized (Pc.+π-cation radical) Derivative. E.N. Bakshi, C.D. Delfs, K.S. Murray, B. Peters, H. Homborg, *Inorg. Chem*, 1988, **27**, 4318.
119. Synthesis and Magnetic Properties of μ-Organaimido-Bridged Iron(III) Salicylaldimine Compounds. Structures of (μ-*p*-Tolylimido)bis[N,N'-ethane-1,2-diylbis(salicylaldiminato) Iron(III), [Fe salen]₂N(tol). P.J. Nichols, G.D. Fallon, K.S. Murray, B.O. West, *Inorg. Chem.*, 1988, **27**, 2795.
118. An Iron-Activated Alcohol Dehydrogenase: Metal Dissociation Constants, Magnetic and Spectroscopic Properties. P.Tse, R.K. Scopes, A.G. Wedd, E. Bakshi, K.S. Murray, *J. Amer. Chem. Soc.*, 1988, **110**, 1295.
117. Applied-field Mössbauer Spectra of Low-Spin Mononuclear and Binuclear Iron(III) Phthalocyanines. E.N. Bakshi, K.S. Murray, *Hyperfine Int.*, 1988, **40**, 283.
116. Synthesis, Structure, ESR Spectra and Redox Properties of (N,N'-ethylenebis)(thiosalicylideneiminato)oxovanadium (IV) and of Related {S,N} Chelates of Vanadium(IV). J.C. Dutton, G.D. Fallon, K.S. Murray, *Inorg. Chem*, 1988, **27**, 34.
115. Binuclear Nickel(II) and Copper(II) Complexes Containing an Endogenous Alkylthiolato-Bridging Sulphur Atom and Exogenous μ-Pyrazolate Group. Crystal Structure of [Ni₂L(pyrazolate)]; (L³⁻ = 1,5-bis(salicylideneiminato)pentane-3-thiolate). P.Iliopoulos, G.D. Fallon, K.S. Murray, *J. Chem. Soc. Dalton Trans.*, 1988, 1823.
114. Synthesis, Reactions and Copper(II), Nickel(II) and Zinc(II) Complexes of N-(2-Pyridinyl)ketothioacetamides. P. Iliopoulos, K.S. Murray, *J. Chem. Soc. Dalton Trans.*, 1988, 433.
113. Molecular and Electronic Structure of Six-Coordinate Low-Spin Oxo-bridged Dimers formed by the Interaction of (μ-oxo)bis(phthalocyaninato) iron(III) with Nitrogen Bases. C. Ercolani, M. Gardini, K.S. Murray, G. Rearesi, B. Rossi, P.R. Zwack, *Inorg. Chem.*, 1987, **26**, 3539.112.

112. Magnetic Properties and Metal-Metal Interactions in Mixed-Metal (Pt₂M) Trimers (M = Ni, Co). D. M. Duckworth, D. M. L. Goodgame, M. A. Hitchman, B. Lippert, K. S. Murray, *Inorg. Chem.* 1987, **26**, 1823-
111. Magnetic properties of high-spin and spin-crossover cobalt(II) Schiff-base compounds. B. J. Kennedy and K. S. Murray, *Inorg. Chim. Acta* 1987, **134**, 249-254.
110. μ -Nitrido-iron-phthalocyanine, (FePc)₂N and of Some Oxidized Derivatives. B.J. Kennedy, K.S. Murray, H. Homborg, W. Kalz, *Inorg. Chim. Acta*, 1987, **134**, 19-21.
109. Zero-Field Splitting and Magnetic Exchange in Chloro(N,N'-ethylenebis(3-methoxysalicylaldehyde)) Iron(III) Hydrate, [Fe(3-MeO-salen)Cl(H₂O)]. B.J. Kennedy, K.S. Murray, *Inorg. Chim. Acta*, 1987, **132**, 153.
108. Structure and magnetism of analogous O-bridged (phenolato) and S-bridged (thiophenolato) binuclear copper(II) complexes. Crystal structure of the binuclear S-bridged complex methanol $(\mu\text{-pyrazolato-N,N'})[\mu\text{-S-a,a-}\{(2\text{-mercapto-4-methylbenzene-1,3-diyl})\text{di(methylidyne)di(azino)}\}\text{bis-(cyclohexanemethanol)ato(3-)}\text{-dicopper(II); L(S)Cu}_2\text{(pt)(CH}_3\text{OH)}$. P. Iliopoulos, K.S. Murray, R. Robson, J. Wilson, G.A. Williams, *J.Chem. Soc., Dalton Trans.*, 1987, 1585.
107. Metal-ligand Bonding Parameters and Magnetic Properties of Some Previously Reported Tetragonal Nickel(II) complexes. B.J. Kennedy, K.S. Murray, M.A. Hitcham, G.L. Rowbottom, *J.Chem. Soc. Dalton Trans.*, 1987, 825.
106. Physical properties of some halogen-doped copper phthalocyanines. B.R. Saunders, K.S. Murray, R.J. Fleming, H. Homborg, *Materials Science* 1987, **XIII**, 231.
105. Structural and electronic properties of six-coordinate Lewis-base adducts of iron(III) Schiff-base complexes. Crystal structure and magnetism of the high-spin complex [(imidazole)Fe(salen)-(NCS)0.5MeOH and of a related complex containing two dissimilar axial ligands. B.J. Kennedy, K.S. Murray, P.R. Zwack, E. Horn, M.R. Snow, E.R.T. Tiekink, *Proc. Indian Acad. Sci. (Chem. Sci.)*, 1987, **98**, 99.
104. Variable temperature magnetic, spectral and X-ray crystallographic studies of 'spin-crossover' iron(III) Schiff-base - Lewis-base adducts. Influence of noncoordinated anions on spin-state interconversion dynamics in [Fe(salen)(imd)₂]Y species (Y = ClO₄⁻, PF₆⁻, BPh₄⁻; imd = imidazole). B.J. Kennedy, A.C. McGrath, K.S. Murray, B.W. Skelton, A.H. White, *Inorg. Chem.*, 1987, **26**, 483.
103. Crystalline isomerism in (μ -oxo)bis(phthalocyaninato)iron(III): Further characterization of the isomer having a linear or quasi-linear Fe-O-Fe bond system (μ -oxo(2)). C. Ercolani, M. Gardini, K.S. Murray, G. Pennesi, G. Ross, *Inorg. Chem.*, 1986, **25**, 3972.
102. Spin states in iron(III)phthalocyanines studied by Mossbauer, magnetic susceptibility and ESR measurements. B.J. Kennedy, K.S. Murray, P.R. Zwack, H. Homborg and W. Kalz, *Inorg. Chem.*, 1986, **25**, 2539.
101. New insights into the electronic structures of solid state dioxygen adducts of N,N'-ethylenebis(salicylaldehyde)cobalt(II), Co(salen) and of active precursors. K.S. Murray, A. van den Bergen, B.J. Kennedy and B.O. West, *Aust. J. Chem.*, 1986, **39**, 1479.

100. Structural and electronic properties of tetranuclear iron(III) pyridyl-amine complexes containing cofacial pairs of μ -oxo-bis(μ -acetato)iron(III) moieties. Models for met hemerythrins. H. Toftlund, K.S. Murray, P.R. Zwack, L.F. Taylor and O.P. Anderson, *J. Chem. Soc., Chem. Commun.*, 1986, 191.
99. Structural, spectroscopic and magnetic properties of mono- and binuclear copper(II) complexes of N-(2-pyridinyl)-ketoacetamides. P. Iliopoulos, G.D. Fallon and K.S. Murray, *J. Chem. Soc., Dalton Trans.*, 1986, 437.
98. Binuclear copper complexes containing hydroxo, phenoxo, alkoxo or thiolato bridging groups in models for a copper binding polymer (SIROREZ)-Cu and related systems, in "Inorganic and Biochemical Aspects of Biological and Inorganic Copper Chemistry", Vol. 2, K.S. Murray, Ed. K.D. Karlin and J. Zubietta, Adenine Press, New York, 1986, pp. 161-178.
97. Magnetostructural correlations in ferromagnetic and antiferromagnetic isomers of a binuclear copper(II) complex. The importance of the geometry of the bridging alkoxo oxygen atom. G.D. Fallon, K.S. Murray, W. Mazurek and M.J. O'Connor, *Inorg. Chim. Acta*, 1985, **96**, L53.
96. Copper thioamide complexes. Synthesis and structures of Di- μ -chlorobis-(N-2-pyridyl)benzenecarbothioamides-S,N'dicopper(I) and of bis(N-2-pyridyl)-benzenecarbothioamide-S,N'copper(I) chloride, complexes containing tetrahedrally coordinated copper centres. M.W. Fuller, V. Costanzo, K.S. Murray, D.St.C. Black, T.W. Hambley and M.R. Snow, *Aust. J.Chem.*, 1985, **38**, 865.
95. μ -oxo-iron(III) phthalocyanine. Electronic structure of the solid form obtained from a dihydroxo-iron(III) precursor. B.J. Kennedy, K.S. Murray, P.R. Zwack, H. Homborg and W. Kalz, *Inorg. Chem.*, 1985, **24**, 3302.
94. Reactions of oxomolybdenum(VI) hydroxamate and thiohydroxamate complexes with hydrazines. Synthesis and structures of a diazenido-hydrazido(1-) complex containing an "end-on"-bonded hydrazido(1-) ligand and of a diazene-diazenido(1-) derivative. M.D. Fitzroy, J.M. Frederiksen, K.S. Murray and M.R. Snow, *Inorg. Chem.*, 1985, **24**, 3265.
93. Magnetic interactions in metal complexes of binucleating ligands. 2. Synthesis and properties of binuclear copper(II) compounds containing exogenous ligands which bridge through two atoms. Crystal and molecular structure of a binuclear- μ -pyrazolato-N:N'-(1)-bridged di-copper(II) complex of 1,3-bis(salicylideneamino)propan-2-ol. W. Mazurek, B.J. Kennedy, K.S. Murray, M.J. O'Connor, J.R. Rodgers, M.R. Snow, A.G. Wedd and P.R. Zwack, *Inorg. Chem.*, 1985, **24**, 3258.
92. Preparation and spectral magnetic and electrochemical characterization of a flexible phenoxo-bridged binuclear copper(II) complex. W. Mazurek, A.M. Bond, K.S. Murray, M.J. O'Connor and A.G. Wedd, *Inorg. Chem.*, 1985, **24**, 2484.
91. Structural and electronic properties of six-coordinate mixed aquo-imidazole iron(III) Schiff-base complexes. Crystal structure of [Fe(3MeO-salen)(5Ph-imd)(H₂O)]BPh₄. B.J. Kennedy, G. Brain, E. Horn, K.S. Murray and M.R. Snow, *Inorg. Chem.*, 1985, **24**, 1647.
90. Oxochromium compounds I. Synthesis and properties of μ -oxo chromium-iron porphyrin and phthalocyanine compounds. D.J. Liston, B.J. Kennedy, K.S. Murray and B.O. West, *Inorg. Chem.*, 1985, **24**, 1561.

89. Magnetic properties and zero-field splitting in high-spin manganese(III) complexes II. Axially ligated manganese(III) porphyrin complexes. B.J. Kennedy and K.S. Murray, *Inorg. Chem.*, 1985, **24**, 1557.
88. Magnetic properties and zero-field splitting in high-spin manganese(III) complexes I. Mononuclear and polynuclear Schiff-base chelates. B.J. Kennedy and K.S. Murray, *Inorg. Chem.*, 1985, **24**, 1552.
87. Iron complexes of N-substituted thiosalicylideneimines III. Structure, magnetism and Mossbauer spectra of chloro(N,N'-ethylenebis(thiosalicylideneiminato))iron(III). An intermediate-spin complex which exists in three crystalline modifications. G.D. Fallon, B.M. Gatehouse, P.J. Marini, K.S. Murray and B.O. West, *J. Chem. Soc. Dalton Trans.*, 1984, 2733.
86. A model for the cyanide form of oxidized cytochrome oxidase: an iron-(III)/copper(II) porphyrin complex displaying ferromagnetic coupling. M.J. Gunter, K.J. Berry and K.S. Murray, *J. Am. Chem. Soc.*, 1984, **106**, 4227.
85. Preparation and characterization of biscyanophthalocyaninatoferrate(III). W. Kalz, H. Homborg, H. Kupfers, B.J. Kennedy and K.S. Murray, *Z. Naturforsch.*, 1984, **39b**, 1478.
84. Crystal structure of a novel μ -hydroxo copper(II) piperazine phenol complex containing a short copper-copper distance and displaying weak exchange coupling. G.D. Fallon, K.S. Murray, B. Spethmann, J.K. Yandell, J.H. Hodgkin and B.C. Loft, *J. Chem. Soc., Chem. Commun.*, 1984, 1561.
83. Preparation and properties of anionic fluoro-complexes of magnesium, zinc and aluminium phthalocyanines, H. Homborg and K.S. Murray, *Z. Anorg. allg. Chem.*, 1984, **517**, 149.
82. Variable deprotonation of an aromatic amino ligand by molybdenum(VI). Crystal and molecular structure of mer-Bis[2,2'-{1,2-phenylenebis(imino-methylene)}-bis(phenolato(3-))]molybdenum(VI). S.F. Gheller, T.W. Hambley, M.R. Snow, K.S. Murray and A.G. Wedd, *Aust. J. Chem.*, 1984, **37**, 911.
81. Chlorophthalocyanine Iron(III). FePc(-2)Cl. A spin admixed ($S=3/2 / S = 5/2$) system. B.J. Kennedy, G. Brain and K.S. Murray, *Inorg. Chim. Acta*, 1984, **81**, L29-L31.
80. Spin-state differences and spin-crossovers in five-coordinate Lewis-base adducts of cobalt(II) Schiff-base complexes. Structure of High-spin [cobalt(II)-saloph-2-methylimidazole adduct. B.J. Kennedy, G.D. Fallon, B.M. Gatehouse and K.S. Murray. *Inorg. Chem.*, 1984, **23**, 580.
79. Unusual structural, chemical and magnetic properties of mononuclear Fe(III) complexes of the potentially binucleating ligand *meso*-tetrakis- $\alpha,\alpha,\alpha,\alpha$ -O-(nicotinamidophenyl)porphyrin. M.J. Gunter, G.M. McLaughlin, K.J. Berry, K.S. Murray, M. Irving and P.E. Clark, *Inorg. Chem.*, 1984, **23**, 283.
78. Structure, magnetism and Mossbauer spectrum of the five-coordinate complex chlorobis (N-methylbenzo-thiohydroxamato)-iron(III). K.J. Berry, P.E. Clark, K.S. Murray, C.L. Raston and A.H. White, *Inorg. Chem.*, 1983, **22**, 3928.
77. Iron complexes of N-substituted thiosalicylideneimines II. Magnetic and Mossbauer spectral studies of iron(II) complexes. P.J. Marini, K.J. Berry, K.S. Murray, B.O. West, M. Irving and P.E. Clark, *J. Chem. (Dalton)*, 1983, 879.

76. Iron complexes of N-substituted Thiosalicylideneimines I. Synthesis and Reactions with Oxygen and Carbon Monoxide. P.J. Marini, K.S. Murray and B.O. West. *J.Chem. Soc. Dalton*, 1983, 143.
75. Specialist periodical reports. Electronic structure and magnetism of inorganic compounds. Vol. 7. Chapter on magnetic susceptibility. P.D.W. Boyd and K.S. Murray, 1982, 134.
74. The formation of heterobinuclear μ -oxo Cr-O-Fe porphyrin complexes by reaction of oxochromium(IV) and Iron(II) porphyrins. D.J. Liston, K.S. Murray and B.O. West, *J.C.S. Chem. Comm.*, 1982, 1109.
73. Magnetic exchange in trans-tetraamminedinitronickel(II), B.N. Figgis, B.J. Kennedy, K.S. Murray, P.A. Reynolds and S. Wright, *Aust. J. Chem.*, 1982, 35, 1807.
72. Magnetic interactions in metal complexes of pentadentate binucleating ligands I. Synthesis and properties of binuclear copper(II) compounds containing single atom bridging ligands. The crystal and molecular structure of a binuclear μ -hydroxo-bridged copper(II) complex of 1,5-bis{1-pyridin-2-yl}ethylideneamino} pentan-3-ol. W. Mazurek, K.J. Berry, K.S. Murray, M.J. O'Connor, M. Snow and A.G. Wedd, *Inorg. Chem.*, 1982, **21**, 3071.
71. Electron spin resonance and electrochemical study of thiohydroxamate and 1-phenyl-3-imino-2(1H)-pyridinethione complexes of copper(II). J. Becher, D.J. Brockway, K.S. Murray, P.J. Newman and H. Toftlund, *Inorg. Chem.*, 1982, **21**, 1791.
70. Water soluble cobalt(II) complexes of N,N'-substituted bis(salicylaldimino-5-sulphonic acids). Oxygen carrying properties and conversion into Co(III) organometallic compounds. K.J. Berry, F. Moya, K.S. Murray, A. van den Bergen and B.O. West, *J. Chem. Soc. (Dalton)*, 1982, 109.
69. Models for functional intermediates in the reaction of cytochrome c oxidase with oxygen, in Proceedings of 2nd BOC/Priestley Conference 'Oxygen and Life' (Royal Society of Chemistry, 1981). K.J. Berry, M.J. Gunter and K.S. Murray, 170.
68. The synthetic model approach to the active site structure of cytochrome oxidase. Novel porphyrin-containing heterobinuclear Fe(III) - Cu(II) complexes. M.J. Gunter, L.N. Mander, K.S. Murray and P.E. Clark, *J. Amer. Chem. Soc.*, 1981, **103**, 6748.
67. Oxygenation reactions of cytochrome oxidase models. Evidence for heme formation in mononuclear and heterobinuclear complexes. M.J. Gunter, L.N. Mander and K.S. Murray, *J.C.S. Chem. Comm.*, 1981, 799.
66. The reaction of N,N'-ethylenebis(mercaptobenzaldiminato)iron(II) with dioxygen. Formation of a μ -peroxo intermediate and isolation of a μ -oxo complex with the unusual $S = 1/2$ spin state. P.J. Marini, K.S. Murray and B.O. West, *J.C.S. Chem. Comm.*, 1981, 726.
65. Electronic properties of thiolate compounds of oxomolybdenum(V) and their tungsten and selenium analogues: effects of ^{17}O , ^{98}Mo and ^{95}Mo -isotope substituents upon esr spectra. G.R. Hanson, A.A. Brunette, A.C. McDonnell, A.G. Wedd and K.S. Murray, *J. Amer. Chem. Soc.*, 1981, **103**, 1953.
64. Specialist periodical reports. Electronic structure and magnetism of inorganic compounds, Vol. 6. Chapter on magnetic susceptibility. P.D.W. Boyd, A.K. Gregson and K.S. Murray, 1980, 233.

63. Synthetic models for cytochrome c oxidase: magnetic and Mössbauer effect studies of binuclear Fe(III) porphyrin-Cu(II) complexes. K.J. Berry, P.E. Clark, M.J. Gunter and K.S. Murray, *Nouveau J. de Chimie*, 1980, **4**, 581.
62. Magnetic exchange in amido-bridged binuclear copper(II) complexes. K.J. Berry, D.St.C. Black, C.H. Bos van der Zalm, K.S. Murray and G.I. Moss, *Inorg. Chim. Acta*, 1980, **46**, 221.
61. Towards synthetic models for cytochrome oxidase: a binuclear Fe(III) porphyrin - Cu(II) complex. M.J. Gunter, L.N. Mander, G.M. McLaughlin, K.J. Berry, P.E. Clark, K.S. Murray and D.A. Buckingham, *J. Amer. Chem. Soc.*, 1980, **102**, 1470.
60. Structure and spectral and redox properties of a dioxomolybdenum(VI) chelate of N-methyl-*p*-tolylthiohydroxamic acid, a complex containing a {O₂MoO₂S₂} donor set. C.A. Cliff, G.D. Fallon, B.M. Gatehouse, K.S. Murray and P.J. Newman, *Inorg. Chem.*, 1980, **19**, 773.
59. An electrochemical study of thiohydroxamate and hydroxamate complexes of iron(III). D. Brockway, K.S. Murray and P.J. Newman, *J. Chem. Soc. (Dalton)*, 1980, 1112.
58. Redox and electron spin resonance properties of oxo-thiolato compounds of molybdenum(V). J. R. Bradbury, G.R. Hanson, I.W. Boyd, S.F. Gheller, A.G. Wedd, K.S. Murray and A.M. Bond. Proceedings of Third Climax Molybdenum Conference, Chicago, 1979.
57. Structure and absolute configuration of tris(N-methylthioformo-hydroxamato)iron(III), an antibacterial metal chelate and potential iron siderochrome. K.S. Murray, P.J. Newman and D. Taylor, *J. Amer. Chem. Soc.*, 1978, **100**, 2251.
56. Mononuclear oxothiolato compounds of molybdenum(V). I.W. Boyd, I.G. Dance, K.S. Murray and A.G. Wedd, *Aust. J. Chem.*, 1978, **31**, 279.
55. Structure, absolute configuration and electrochemical properties of tris-N-methylformothiohydroxamato)iron(III), an antibacterial metal chelate and potential iron siderochrome. K.S. Murray, P.J. Newman, B.M. Gatehouse and D. Taylor, *Aust. J. Chem.*, 1978, **31**, 983.
54. Ruthenium complexes with a tetradentate salicylaldimine Schiff-base. A. van den Bergen, K.S. Murray and B.O. West, *Aust. J. Chem.*, 1978, **31**, 203.
53. Synthesis and electron spin resonance study of a tetradentate copper(II) mercaptobenzaldimine complex. M.F. Corrigan, K.S. Murray, B.O. West and J.R. Pilbrow, *Aust. J. Chem.*, 1977, **30**, 2455.
52. Thiohydroxamato complexes of iron(III). Synthesis, magnetism and Mössbauer spectra. A.J. Mitchell, K.S. Murray, P.J. Newman and P.E. Clark, *Aust. J. Chem.*, 1977, **30**, 2439.
51. Cobalt(III) and iron(III) complexes of 2-aminocyclopentene-1-dithio-carboxylic acid. K.S. Murray, M.F. MacKay, M.J. O'Connor, P.J. Oliver and P.J. Newman, *Inorg. Nucl. Chem. Letters*, 1977, **13**, 417.
50. An electron spin resonance study of the oxygenation of a tetradentate thiosalicylideneimine cobalt(II) chelate. M.F. Corrigan, K.S. Murray, B.O. West, P.R. Hicks and J.R. Pilbrow, *J. Chem. Soc. (Dalton)*, 1977, 1478.

49. Models for the bacterial iron transport chelate enterochelin. B.F. Anderson, D.A. Buckingham, G.B. Robertson, J. Webb, P.E. Clark and K.S. Murray, *Nature*, 1976, **262**, 722.
48. Preparation and metal complexes of 2,2'-pyridylbis(2"-pyridyl)hydrazone. Dt.St.C. Black, A.J. Hartshorn and K.S. Murray, *Aust. J. Chem.*, 1976, **29**, 153.
47. Paramagnetic anisotropy and electronic structure of NN'-ethylenebis-(acetylacetoniminato)cobalt(II) and NN'-ethylenebis(thioacetylaceton-iminato)cobalt(II). K.S. Murray and R.M. Sheahan, *J. Chem. Soc. (Dalton)*, 1976, 1134.
46. Paramagnetic anisotropy and electronic structure of N,N'-ethylenebis-(salicylideneiminato)cobalt(II), its pyridine adduct, and N,N'-ethylenebis(thiosalicylideneiminato)cobalt(II). K.S. Murray and R.M. Sheahan, *J. Chem. Soc. (Dalton)*, 1976, 999.
45. Paramagnetic anisotropy and electronic structure of copper(II) tetraphenylporphyrin. K.S. Murray and R.M. Sheahan, *Aust. J. Chem.*, 1975, **28**, 2623.
44. The X-ray structure and magnetic anisotropy of N,N'-ethylenebis(thio-salicylaldiminato)cobalt(II): Co S-Salen. M.F. Corrigan, K.S. Murray, R.M. Sheahan, B.O. West, G.D. Fallon and B.M. Gatehouse, *Inorg. Nucl. Chem. Letters*, 1975, **11**, 625.
43. Ground-state of low-spin cobalt(II) chelate complexes. K.S. Murray and R.M. Sheahan, *Chem. Comm.*, 1975, 476.
42. ¹³C n.m.r. spectra of binuclear iron(III) Schiff-base complexes. K.S. Murray, *J. Chem. Soc. (Dalton)*, 1975, 1538.
41. The synthesis and the magnetic properties of perfluoroalkyl organometallic chromium(III) complexes containing N-substituted salicylaldimines and other ligands. A. van den Bergen, K.S. Murray, R.M. Sheahan and B.O. West, *J. Organomet. Chem.*, 1975, **90**, 299.
40. Amino acid complexes of iron. I. Iron(II) complexes of thioglycollic acid, I-cysteine and related ligands. K.S. Murray and P.J. Newman, *Aust. J. Chem.*, 1975, **28**, 773.
39. Crystal structure of mer-tris(N-benzylsalicylaldiminato)manganese(III). K.S. Murray, C.L. Raston, A.H. White and A.C. Willis, *J. Chem. Soc. (Dalton)*, 1974, 1793.
38. The coordinating properties of N-substituted thiosalicylaldimines with cobalt(II), and cobalt(III) organometallics containing an S₂N₂ quadridentate Schiff-base. A. van den Bergen, M. F. Corrigan, K.S. Murray, R.M. Slade and B.O. West, *Inorg. Nucl. Chem. Letters*, 1974, **10**, 859.
37. Binuclear oxo-bridged iron(III) complexes. K. S. Murray, *Coord. Chem. Revs.*, 1974, **12**, 1.
36. Single-crystal magnetic studies of cobalt(II) Schiff-base chelate compounds. K. S. Murray and R.M. Sheahan, *Chem. Phys. Letters*, 1973, **22**, 406.
35. Crystal and molecular structure and magnetic properties of μ -acetato N,N'-ethylenebis(salicylaldiminato)manganese(III): A linear chain compound containing a single anti-anti acetate bridge. J.E. Davies, B.M. Gatehouse and K.S. Murray, *J. Chem. Soc. (Dalton)*, 1973, 2523.

34. Anionic dimethylplatinum(IV) Schiff-base chelate complexes. K.S. Murray, B.E. Reichert and B.O. West, *J. Organomet. Chem.*, 1973, **63**, 461.
33. Dimethyl gold(III) complexes of salicylaldimine Schiff-bases. K.S. Murray, B.E. Reichert and B.O. West, *J. Organomet. Chem.*, 1973, **61**, 451.
32. Vanadium(III) chloro- and bromo- complexes with tetradentate Schiff-base ligands. K.S. Murray, G.R. Simm and B.O. West, *Aust. J. Chem.*, 1973, **26**, 991.
31. Ferromagnetic coupling in nickel(II) monothiocarbamate complexes. E.M. Krankovits, R.J. Magee, K.S. Murray and M.J. O'Connor, *Aust. J. Chem.*, 1973, **26**, 749.
30. Mixed valence complexes of vanadium with phenanthroline and bipyridyl. K.S. Murray and R.M. Sheahan, *J. Chem. Soc. (Dalton)*, 1973, 1182.
29. Preparation and reduction of Rh(III) complexes of a quadridentate Schiff-base ligand, and synthesis of organo-Rh(III) derivatives. R.J. Cozens, K.S. Murray and B.O. West, *J. Organomet. Chem.*, 1972, **38**, 391.
28. Mixed Schiff-base β -diketone complexes of cobalt(III). R.J. Cozens and K.S. Murray, *Aust. J. Chem.*, 1972, **25**, 711.
27. Molybdenum(IV) halide complexes with the chelating ligands Schiff-bases, acetylacetonone and 8-hydroxyquinoline. A. van den Bergen, K.S. Murray and B.O. West, *Aust. J. Chem.*, 1972, **25**, 705.
26. Magnetic properties of some tridentate Schiff-base manganese(II) complexes. K.D. Butler, K.S. Murray and B.O. West, *Aust. J. Chem.*, 1971, **24**, 2249.
25. Fluoroalkyl-cobalt(III) Schiff-base complexes. A. van den Bergen, K.S. Murray and B.O. West, *J. Organomet. Chem.*, 1971, **33**, 89.
24. Transition metal complexes of thiabendazole. C. Kowala, K.S. Murray, J.M. Swan and B.O. West, *Aust. J. Chem.*, 1971, **24**, 13, 1369.
23. Proton Knight shifts and antiferromagnetic exchange in dimeric oxygen-bridged Schiff-base iron(III) complexes. P.D.W. Boyd and K.S. Murray, *J. Chem. Soc. (A)*, 1971, 2711.
22. Magnetic properties of polynuclear hydroxy-iron(III) sulphate complexes. R.W. Cattrall, K.S. Murray and K.I. Peverill, *Inorg. Chem.*, 1971, **10**, 1301.
21. Rhodium(I) and iridium(I) carbonyl and olefin derivatives of salicylaldimine Schiff-bases. R.J. Cozens, K.S. Murray and B.O. West, *J. Organomet. Chem.*, 1971, **27**, 399.
20. Some organometallic and 'simple' Schiff-base chelates of tin(IV). A. van den Bergen, R.J. Cozens and K.S. Murray, *J. Chem. Soc. (A)*, 1970, 3060.
19. Rhodium(III) - Alkyls stabilized by a quadridentate Schiff-base ligand. R.J. Cozens, K.S. Murray and B.O. West, *Chem. Comm.*, 1970, 1262.
18. Mössbauer effect studies of some dimeric iron compounds. A.N. Buckley, I.R. Herbert, B.D. Rumbold, G.V.H. Wilson and K.S. Murray, *J. Phys. Chem. Solids*, 1970, **31**, 1423.
17. Mössbauer effect and susceptibility studies of a dimeric Schiff-base iron(III) adduct. A.N. Buckley, B.D. Rumbold, G.V.H. Wilson and K.S. Murray, *J. Chem. Soc. (A)*, 1970, 2298.

16. Electron spin resonance study of exchange and dipole-dipole coupling in copper(II) chelates of cyclopentane-tetracarboxylic acid and related systems. J.H. Price, J.R. Pilbrow, K.S. Murray and T.D. Smith, *J. Chem. Soc.(A)*, 1970, 968.
15. Thallium(I) Schiff-base compounds. R.J. Cozens, B.O. West and K.S. Murray, *Aust. J. Chem.*, 1970, **23**, 683.
14. S-sulphinates of cobalt(III) planar chelates and sulphur dioxide insertion into a cobalt(III)-carbon bond. R.J. Cozens, G.B. Deacon, P.W. Felder, K.S. Murray and B.O. West, *Aust. J. Chem.*, 1970, **23**, 481.
13. Electronic spin relaxation in dimeric iron(III) complexes. A.N. Buckley, G.V.H. Wilson and K.S. Murray, *Chem. Comm.*, 1969, 718.
12. Relaxation effects in the Mössbauer spectra of some dimeric iron compounds. A.N. Buckley, G.V.H. Wilson, and K.S. Murray, *Solid State Comm.*, 1969, **7**, 471.
11. Magnetic properties and Mössbauer spectra of some terdentate Schiff-base iron(III) halide complexes. A. van den Bergen, K.S. Murray, B.O. West and A.N. Buckley, *J. Chem. Soc., (A)*, 1969, 2051.
10. N-substituted salicylaldimine complexes of manganese(III). Synthesis, magnetism and spectra. A. van den Bergen, K.S. Murray, M.J. O'Connor and B.O. West, *Aust. J. Chem.*, 1969, **22**, 39.
9. Insertion of SO₂ into a cobalt(III)-carbon bond. K.S. Murray, R.J. Cozens, G.B. Deacon, P.W. Felder and B.O. West, *Inorg. Nucl. Chem. Letters*, 1968, **4**, 705.
8. Some new mono and binuclear complexes of iron(III) with tridentate Schiff-bases. K.S. Murray and B.O. West, *Inorg. Nucl. Chem. Letters*, 1968, **4**, 439.
7. N-substituted salicylaldimine complexes of iron(III). (II) Magnetic properties. A. van den Bergen, K.S. Murray and B.O. West, *Aust. J. Chem.*, 1968, **21**, 1517.
6. N-substituted salicylaldimine complexes of iron(III). (I) Synthesis, infrared and mass spectra. A. van den Bergen, K.S. Murray, M.J. O'Connor, Nadja Rehak and B.O. West, *Aust. J. Chem.*, 1968, **21**, 1505
5. Complexes of Fe(III) with N-substituted salicylideneimines. K.S. Murray, A. van den berges, M.J. O'Connor, N. Rehak and B.O. West, *Inorg. Nucl. Chem. Letters*, 1968, **4**, 87.
4. The magnetic properties of some titanium(III) complexes. D.J. Machin, K.S. Murray and R.A. Walton, *J. Chem. Soc.(A)*, 1968, 195.
3. Magnetic and spectral studies on tungsten(V) chloride and bromide and several oxyhalide and halide complexes of molybdenum(V) and tungsten(V). B.J. Brisdon, D.A. Edwards, D.J. Machin, K.S. Murray and R.A. Walton, *J. Chem. Soc.(A)*, 1967, 1825.
2. The magnetism and spectra of some octahedral vanadium(III) complexes. D.J. Machin and K.S. Murray, *J. Chem. Soc.(A)*, 1967, 1498.
1. The magnetic properties of some vanadium(IV) complexes. D.J. Machin, and K.S. Murray, *J. Chem. Soc.(A)*, 1967, 1330.